- PEAKALL, R. and P. E. SMOUSE (2006): Genalex 6: genetic analysis in Excel. Population genetic software for teaching and research. *Mol. Ecol. Notes* **6**: 288–295.
- POSNETTE, A. F. and J. M. A. TODD (1951): Virus diseases of cacao in West Africa. VIII. The search for virus resistant cacao. *Annals of Applied Biology* **38**: 785–800.
- POWELL, W., M. MORGANTE, C. ANDRE, M. HANATEY, J. VOGEL, S. TINGEY and A. RAFALSKI (1996): The comparison of RFLP, RAPD, AFLP and SSR (microsatellite) markers for germplasm analysis. *Molecular Breeding* **2**: 225–238.
- PRITCHARD, J. K., M. STEPHENS and P. DONNELLY (2000): Inference of population structure from multilocus genotype data. *Genetics* **155**: 945–959.
- PUGH, T., O. FOUEY, A. M. RISTERUCCI, P. BROTTIER, M. ABOULADZE, C. DELETREZ, B. COURTOIS, D. CLEMENT, P. LARMANDE, J. A. K. NGORAN and C. LANAUD (2004): A new cacao linkage map based on codominant markers: development and integration of 201 new microsatellite markers *Theor. Appl. Genet* **108**: 1151–1161.
- PURSEGLOVE, J. W. (1974): Tropical crops, dicotyledons. John Wiley & Sons, New York.
- QUENOILLE, M. (1956): Notes on bias in estimation. *Biometrika* **43**: 253–260.
- RICE, W. R. (1989): Analyzing tables of statistical tests. *Evolution* **43**: 223–225.
- RICE, R. A. and R. GREENBERG (2000): Cacao cultivation and the conservation of biological diversity. *Ambio* **29**(3): 167–173.
- SAUNDERS, J. A., S. MISCHKE, E. A. LEAMY and A. A. HEMEIDA (2004): Selection of international molecular standard for DNA fingerprinting. *Theor. Appl. Genet.* **110**: 41–47.
- SCHNELL, R. J., C. T. OLANO, J. S. BROWN, A. W. MEEROW, C. CERVANTES-MARTINEZ, C. NAGAI and J.-C. MOTAMAYOR (2005): Retrospective determination of the parental population of superior cacao (*Theobroma cacao* L.) seedlings and association of microsatellite alleles with productivity. *J. Amer. Soc. Hort. Sci.* **130**: 181–190.
- SERENO, M., P. P. ALBUQUERQUE, R. VENCovsky and A. FIGUERA (2006): Genetic diversity and natural population structure of cacao, *Theobroma cacao* L. from the Brazilian Amazon evaluated by microsatellite markers. *Conservation Genetics* **7**: 13–24.
- SWEIGART, A., K. KAROLY, A. JONES and J. H. WILLIS (1999): The distribution of individual inbreeding coefficients and an pairwise relatedness in a population of *Mimulus guttatus*. *Heredity* **83**: 625–632.
- TOXOPEUS, H. (1964): F₃ Amazon cacao in Nigeria. *Rep. Cacao Res. Inst. Nigeria*, 1963/64, pp.13–22.
- WEIR, B. S. and C. C. COCKERHAM (1984): Estimating F-statistics for the analysis of population structure. *Evolution* **38**: 1358–1370.
- WRIGHT, S. (1965): The interpretation of genetic structure by F-statistics with special regard to systems of mating. *Evolution* **19**: 355.
- ZHANG, D., E. AREVALO, B. MISCHKE, L. ZUNIGA and A. BARRETO (2006a): Genetic diversity and population structure of cocoa, *Theobroma cacao* L. in the Hulla and Ucayali valleys of Peru. *Annals of Botany* **98**: 647–655.
- ZHANG, D., E. AREVALO, S. MISCHKE, R. GOENAGA, A. A. HEMEIDA and J. A. SAUNDERS (2006b): Accuracy and reliability of high-throughput microsatellite genotyping for cacao clone identification. *Crop Science* **46**: 2084–2092.
- ZHANG, D., M. BOCCARA, L. MOTILAL, D. R. BUTLER, P. UMAHARAN, S. MISCHKE and L. MEINHARDT (2008): Microsatellite variation and population structure in the “refractorio” cacao of Ecuador. *Conserv. Genet* **9**: 327–337.
- ZHANG, D., S. MISCHKE, E. S. JOHNSON, W. PHILLIPS-MORA and L. MEINHARDT (2009): Molecular characterization of an International cocoa collection using microsatellite markers. *Tree Genetics & Genomes* **5**: 1–10.

Cross-amplification and Characterization of Polymorphic Microsatellite Markers From *Acacia* (Senegalia) *mellifera* and *Acacia brevispica* to *Acacia senegal* (L.) Willd.

By S. F. OMONDI^{1,3,*}, O. G. DANGASUK², D. W. ODEE^{3,4}, S. CAVERS⁴ and D. P. KHASA⁵

(Received 6th May 2009)

¹) Department of Forestry and Wood Science, Moi University, P.O. Box 1125 Eldoret, Kenya.

²) Department of Biological sciences, Moi University, P.O. Box 1125 Eldoret, Kenya.

³) Kenya Forestry Research Institute, P.O. Box 20412-00200 Nairobi, Kenya.

⁴) Centre for Ecology and Hydrology, Bush Estate, Penicuik, Midlothian, EH26 0QB, UK.

⁵) Centre for Forest Research and Canada Research Chair in Forest and Environmental Genomics, Université Laval, Sainte-Foy, Québec, Canada G1V 0A6.

* Author of correspondence: STEPHEN F. OMONDI.
E-Mail: stephenf.omondi@gmail.com

Abstract

Seven polymorphic microsatellite markers isolated from *Acacia brevispica* and *Acacia mellifera* were successfully cross-amplified in *Acacia senegal*. The loci were surveyed for polymorphism using 30 samples. Allelic diversity ranged from 4 (*Ame02*, *Ab06* and *Ab18*) to 13 (*Ab26*) per locus. The expected heterozygosity (H_e) ranged from 0.543 (*Ame02*) to 0.868 (*Ab26*) while observed heterozygosity (H_o) ranged from 0.516 (*Ame05*) to 0.800 (*Ame03*). Cross-amplification of these loci represents a potential source of co-dominant markers and will be useful in the study of genetic diversity, structure,

gene flow and breeding systems of this important *Acacia* species.

Key words: *Acacia senegal*, cross-amplification, genetic diversity, microsatellite, Sub-Saharan Africa.

Introduction

Acacia senegal (L.) Willd. is a multipurpose agroforestry tree, native to the arid and semi-arid lands of Sub-Saharan Africa but found as far as the Indian sub-continent. It grows to a height of 5–12 m, with a trunk up to 30 cm in diameter (FAGG and ALLISON, 2004) and is valued for gum arabic production, as well as secondary

agricultural roles, such as restoring soil fertility and providing fuel and fodder (BALLAL *et al.*, 2005). It is also essential in sand dune fixation and is preferred in bush-fallow rotation and intercropping (ANDERSON and WEI-PING, 1992). The species has four varieties: *senegal*, *kerensis*, *rostrata* and *leiorhachis* but their delimitation is still problematic (FAGG and ALLISON, 2004). Intraspecific variation has been reported in morphological, molecular and biochemical characteristics (CHIKAMAI and ODERA, 2002; MOTLAGH *et al.*, 2006; AL-ASSAF *et al.*, 2007). Despite its ecological and economic importance, little progress has been made in developing sets of molecular markers for genetic studies.

Table S1. – Microsatellite loci isolated for *A. brevispica* (SCHNABEL *et al.*, 2005) and *A. mellifera* (RUIZ-GUAJARDO *et al.*, 2007) used in cross-amplification study in *A. senegal*.

Locus Accession no.	Microsatellite repeat	Loci amplification
Ab01	(AAG) ₁₅	Irregular amplification
Ab02	(AC) ₁₁	Poor amplification
Ab03	(AC) ₁₇	No amplification
Ab04	(AC) ₇	No amplification
Ab05	(AG) ₈ ...	No amplification
Ab06	(AC) ₉ (AG) ₁₀	Polymorphic amplification
Ab07	(ATC) ₇	Irregular amplification
Ab09	(AAAC) ₆	No amplification
Ab10	(AG) ₁₃	No amplification
Ab12	(AG) ₁₁	Irregular amplification
Ab13	(AG) ₁₉	No amplification
Ab14	(AG) ₁₀	Irregular amplification
Ab15	(AC) ₁₀	Poor amplification
Ab16	(AT) ₈ (AG) ₁₁	Irregular amplification
Ab17	(AAG) ₁₄ (ATC) ₁₁	Poor amplification
Ab18	(AAG) ₁₅	Polymorphic amplification
Ab19	(AC) ₁₆	No amplification
Ab20	(AG) ₂₄ (AC) ₁₆	No amplification
Ab21	(AGAT) ₂₃	No amplification
Ab22	(AG) ₁₁	Poor amplification
Ab23	(AC) ₉ (AC) ₇	No amplification
Ab25	(AC) ₇	Poor amplification
Ab26	(AG) ₈ (AG) ₉	Polymorphic amplification
Ab27	(AC) ₇	No amplification
Ame01	(AAG) ₇	No amplification
Ame02	(AC) ₇ (AG) ₁₁	Polymorphic amplification
Ame03	(AG) ₉	Polymorphic amplification
Ame04	(AG) ₁₀	Poor amplification
Ame05	(ATC) ₇	Polymorphic amplification
Ame06	(GT) ₁₀	Poor amplification
Ame07	(GT) ₂₀	Polymorphic amplification
Ame08	(GTT) ₉	No amplification
Ame09	(GA) ₁₇	No amplification
Ame10	(CA) ₉	No amplification
Ame12	(CT) ₁₂ (AC) ₆	No amplification

S1- Supplementary material (*Table 1*).

Table 1. – Microsatellite loci in *A. senegal* obtained from cross amplification from *A. mellifera* and *A. brevispica* including GenBank accession number; Ta – annealing temperature (touchdown range); bp – expected allelic size in base pairs; A – number of alleles observed per locus; n – number of individuals used; x, number of alleles recorded for the original species; H_O – observed heterozygosity; y, H_O recorded for the original species; H_E – expected heterozygosity; z, H_E recorded for the original species.

Accession no.	Primer sequence	Repeat motif	Ta (°C)	Size (bp)	A (n) {x}	H _O {y}	H _E {z}
<i>Ame02</i>	F-GAACCATCAGCGTAATAA	(AC) ₇ (AG) ₁₁	55-45	117	4 (30) {7}	0.670 {0.792}	0.543 {0.802}
<i>DQ467674</i>	R-GGTTTAGCAACATACTATCTC						
<i>Ame03</i>	F-GAACAATATCAGCAATCACT	(AG) ₉	55-45	139	8 (30) {4}	0.800 {0.271}	0.810 {0.339}
<i>DQ467673</i>	R-CCTCATGCACACACAAGAT						
<i>Ame05</i>	F-CCCAACAAAGATCATCAT	(ATC) ₇	58-48	203	5 (30) {13}	0.516 {0.532}	0.530 {0.580}
<i>DQ467656</i>	R-ATGGTTCAGTTTCTTTATTCT						
<i>Ame07</i>	F-ATAAAAAACAAAACCCAACTAAATG	(GT) ₂₀	55-45	353	10 (30) {19}	0.695 {0.958}	0.802 {0.911}
<i>DQ467658</i>	R-GTCCAAACTCTTCAATGTCAA						
<i>Ab06</i>	F-CCTTCTTTGACGGTATTC	(AC) ₉ (AG) ₁₀	58-48	147	4 (30) {7}	0.785 {1.00}	0.745 {0.911}
<i>AY843537</i>	R-TCATCTCTCTCTCCATT						
<i>Ab18</i>	F-GAAGGGTCTGGCATTAC	(AAG) ₁₅	60-50	212	4 (30) {3}	0.717 {0.312}	0.579 {0.280}
<i>AY843549</i>	R-CGACGACGAAGATACT						
<i>Ab26</i>	F-ATATTCTGCTTAGTCTA	(AG) ₈ (AG) ₉	61-51	126	13 (30) {6}	0.780 {0.687}	0.868 {0.752}
<i>AY843557</i>	R-GGGGCATAAATATGAG						

* All the forward primers were tagged with M13R tag 5'-CACGACGTTGTAAAACGAC-3'.

Microsatellites are among the best classes of markers for studying genetic processes in natural populations because they are abundant, co-dominant, interspersed throughout the genome, possess high resolution power and polymorphism (CHASE *et al.*, 1996). Their development, however, requires a significant investment (FERNANDEZ *et al.*, 2000); therefore, when markers have been developed in taxonomically related species, it is worthwhile testing for cross-amplification. In this brief preliminary study thirty-four microsatellite primers developed from *A. mellifera* and *A. brevispica* were tested for cross-amplification in *A. senegal*.

Materials and Methods

Microsatellite loci screening

Genomic DNA was isolated from leaves following a modified CTAB procedure (FERNANDEZ *et al.*, 2000). Thirty four microsatellite loci isolated from *Acacia mellifera* (RUIZ-GUAJARDO *et al.*, 2007) and *Acacia brevispica* (SCHNABEL *et al.*, 2005) were tested for amplification in 4 *A. senegal* DNA samples. PCR runs containing 20 ng of DNA, 1x PCR buffer (10 mM Tris-HCL (pH 8.3), 50 mM KCl), 2.5 mM MgCl₂, 200 µM of each dNTP, 0.1 µM of each primer, and 1 unit of *Taq* DNA polymerase (Invitrogen), were performed in a volume of 10 µl (RUIZ-GUAJARDO *et al.*, 2007, OTERO-ARNAIZ *et al.*, 2005). Reactions were carried out on an MJ Research thermal cycler following a touchdown protocol: 95°C for 3 minutes; 20 cycles of 95°C for 30 seconds, either 60°C–50°C, 58°C–48°C or 55°C–45°C (–0.5°C) for 30 seconds, 72°C for 30 seconds; 10 cycles of 95°C for 30 seconds, 50°C, 48°C or 45°C for 30 seconds, 72°C for 30 seconds; final extension of 72°C for 10 minutes and held at 4°C. PCR products were electrophoresed alongside 100 bp ladder, stained in ethidium bromide and visualized under UV-light. Loci that produced single, distinct bands at the expected size were selected for further screening.

Forward primers of the selected loci were tagged with M₁₃ fluorescent tail and optimized using different annealing temperatures, PCR buffers, Mg²⁺, dNTPs, primer and *Taq* DNA polymerase concentrations. The mix that produced sharpest and most distinct bands at the expected size was selected (20 ng of genomic DNA, 1 X PCR buffer (10 mM Tris-HCl pH 8.3, 50 mM KCl), 1.5 mM MgCl₂, 200 µM of each dNTP, 0.05 µM of each primer, and 1 unit of *Taq* DNA polymerase). These loci were then tested for polymorphism and reproducibility.

Polymorphism testing

The final set of loci was screened for polymorphism using a set of 30 DNA samples (3 samples per population from 10 populations). PCR was carried out as described above and products separated on 0.25 mm denaturing polyacrylamide gels containing 10% Ammonium Persulfate (APS), tetramethylethylenediamine (TEMED)-Omnipur and 8% acrylamide (Gene-PAGE plus)-Amresco and electrophoresed using 1 X TBE buffer on a Li-Cor IR² 4200 DNA sequencer. Band sizes were determined by referencing standard IRD800, 50–350 bp and 50–700 bp ladder (KHASHA *et al.*, 2005). Loci that were polymorphic and informative were then selected as a set of *A. senegal* markers for further studies.

Results

Out of the 34 loci screened, 11 were selected for polymorphism screening with a larger number of DNA samples while others failed to amplify or showed poor amplification (Table S1). During the screening, 7 loci (Table 1) showed positive amplification and unambiguous patterns of polymorphic bands and were selected as *A. senegal* markers. The loci showed features desired for co-dominant molecular markers in the species. No deviation from Hardy Weinberg equilibrium or linkage dise-

equilibrium was detected between loci ($P > 0.05$) (RAYMOND and ROUSSET, 1995). The number of alleles and expected heterozygosity were also estimated for each locus. The total number of alleles was highly variable across loci, ranging from 4 (*Ame02*, *Ab06* and *Ab18*) to 13 (*Ab26*) (*Table 1*). Expected heterozygosity ranged from 0.543 (*Ame02*) to 0.868 (*Ab26*) (*Table 1*). Overall, the loci proved to be informative, at levels comparable to the source species and microsatellite loci of other tropical tree species.

Conclusion

This study unveiled evidence that cross-transferability of developed microsatellite loci can increase the availability of markers to address both ecological and evolutionary questions in *A. senegal*. These loci will form a useful marker set for population genetic studies of the species. These will include detecting and understanding available genetic diversity within the species range for maximum utilization and conservation purposes.

Acknowledgements

This work was funded by Department of Foreign Affairs and International Trade Canada (DFAIT) through Canadian Bureau for International Education (CBIE) in the framework of Graduate Student Exchange program (GSEP) grant for 2007 and Kenya Forestry Research Institute (KEFRI), through ACACIAGUM project. We thank Mr. JOHN GICHERU of KERFI and ANDRÉ GAGNÉ of Center for Forest Research, Université Laval for coordinating the acquisition of research materials and all colleagues at the bioinformatics Centre for their advice and support.

References

- AL-ASSAF, S., G. O. PHILLIPS, H. AOKI and Y. SASAKI (2007): Characterization and properties of *Acacia senegal* (L.) Willd. var. *senegal* with enhanced properties (*Acacia* (sen) SUPER GUM™): Part 1. Controlled maturation of *Acacia senegal* var. *senegal* to increase viscoelasticity, produce a hydrogel form and convert a poor into a good emulsifier. *Food Hydrocolloids* **21**: 319–328.
- ANDERSON, D. M. W. and P. W. WEIPING (1992): Gum arabic (*Acacia senegal*) from Uganda: characteristics NMR spectra, amino compositions and gum/soil cationic relationships. *International Tree Crops Journal* **7**: (3), 167–179.
- BALLAL, M. E., E. A. EL-SIDDIG, M. A. ELFADL and O. LUUKKANEN (2005): Relationship between environmental factors, tapping dates, tapping intensity and gum arabic yield of an *Acacia senegal* plantation in western Sudan. *Journal of Arid Environments* **63**: 379–389.
- CHASE, M., R. KESSELI and K. BAWA (1996): Microsatellite markers for population and conservation genetics of tropical trees. *American Journal of Botany* **83**: (1), 51–57.
- CHIKAMAI, B. N. and J. A. ODERA (2002): Commercial plant gums and resins in Kenya. Nairobi, Kenya.
- FAGG, C. W. and G. E. ALLISON (2004): *Acacia senegal* and Gum Arabic Trade. Oxford Forestry Institute, Tropical Forestry Papers. No. 42.
- FERNÁNDEZ, J. F., V. L. SORK, G. GALLEGÓ, J. LÓPEZ, A. BOHORQUES and J. TOHME (2000): Cross-Amplification of microsatellite loci in a Neotropical *Quercus* Species and standardization of DNA extraction from mature leaves Dried in silica gel. *Plant Molecular Biology Reporter* **18**: 397–397.
- KHASA, P. D., P. POLLEFEYS, A. NAVARRO-QUEZADA, P. PERINET and J. BOUSQUET (2005): Species-specific Microsatellite markers to monitor gene flow between exotic poplars and their natural relatives in eastern North America. *Molecular Ecology Notes* **5**: 920–923.
- MOTLAGH, S., P. RAVINES, K. A. KARAMALLAH and M. QIFENG (2006): The analysis of *Acacia* gums using electrophoresis. *Food Hydrocolloids* **20**: (6), 848–854.
- RAYMOND, M. and F. ROUSSET (1995): GENEPOP Version 1.2: population genetics software for exact tests and ecumenicism. *Journal of Heredity* **86**: 248–249.
- RUIZ-GUAJARDO, J. C., A. OTERO-ARNAIZ, T. TAYLOR, G. STONE, C. T. GLENN, N. A. SCHNABEL, J. T. MILLER, S. PREUSS and A. SCHNABEL (2007): Isolation of polymorphic microsatellite markers in the sub-Saharan tree, *Acacia* (senegalia) *mellifera* (Fabaceae: Mimosoideae). *Molecular Ecology Notes* **7**: 1138–1140.
- SCHNABEL, A., A. OTERO-ARNAIZ, T. C. GLENN, N. A. SCHNABEL, C. HAGENT and L. NDONG (2005): Isolation and characterization of microsatellite markers in the East African tree, *Acacia brevispica* (Fabaceae: Mimosoideae). *Molecular ecology notes* **5**: 366–368.