Acta Scientifica Naturalis

Former Annual of Konstantin Preslavsky University of Shumen: Chemistry, Physics, Biology, Geography

Journal homepage: asn.shu.bg

Chemical and geochemical modeling. Thermodynamic models for binary fluoride systems from low to very high concentration (> 35 m) at 298.15 K

Stanislav Donchev, Tsvetan V. Tsenov, Christomir Christov

University of Shumen Bishop Konstantin Preslavski, Department of Chemistry, Faculty of Natural Sciences,
115 Universitetska Str., 9700 Shumen, Bulgaria

Abstract: *In this study we developed well validated thermodynamic models for solution behavior and solid-liquid equilibrium for all fluoride binary systems, for which activity data are available. The subject of modeling study are 5 fluoride systems of the type 1-1 (HF-H₂O, NaF-H₂O, KF-H₂O, RbF-H₂O, and CsF-H₂O) and one of 1-2 type (H₂SiF₆-H₂O) from low to very high concentration at 298.15 K. Models are developed on the basis of Pitzer ion interactions approach. The recommendations on mean activity coefficients ($\gamma_{\pm}$) have been used to construct the model for HF-H₂O system. To parameterize models for all other 5 binary systems we used all available raw experimental osmotic coefficients data (ϕ) for whole concentration range of solutions, and up to saturation point. The predictions of new developed here models are in excellent agreement with experimental osmotic coefficients data, and with recommendations on activity coefficients in binary solutions from low to very high concentration: up to 20 mol. kg⁻¹ in HF-H₂O, and up to 35.6 mol.kg⁻¹ in CsF-H₂O. The Deliquescence Relative Humidity (DRH (%)) and thermodynamic solubility products (as $\ln K_{sp}^{\circ}$) of 4 solid phases [NaF(s), KF.2H₂O(s), RbF(s), and CsF(s)] have been determined on the basis of evaluated model parameters and using experimental $m(\text{sat})$ solubility data.*

Keywords: Computer chemical modeling; Pitzer approach; Binary fluoride systems (HF-H₂O, H₂SiF₆-H₂O, NaF-H₂O, KF-H₂O, RbF-H₂O, and CsF-H₂O); Solution activity; Solid-liquid phase equilibrium

Introduction

Computer models that predict solution behavior and solid-liquid-gas equilibria close to experimental accuracy have wide applicability. They can simulate the complex changes that occur in nature and can replicate conditions that are difficult or expensive to duplicate in the laboratory. Such models can be powerful predictive and interpretive tools to study the geochemistry of natural waters and mineral deposits, solve environmental problems and optimize industrial processes. However, development of comprehensive not low- and high- concentration restricted models is a very difficult, time consuming and challenging task. The specific interaction approach for describing electrolyte solutions to high concentration introduced by Pitzer (1973, 1991) represents a significant advance in physical chemistry that has facilitated the construction of accurate computer thermodynamic models. It was showed that this approach could be expanded to accurately calculate solubilities in complex brines, and to predict the behavior of natural and industrial fluids from very low to very high concentration at standard temperature of 25°C (Harvie et al., 1984; Andre et al., 2019; Lach et al., 2018; Guignot et al., 2019; Park et al. 2009; Kolev et al., 2013; Lassin et al., 2020; Christov, 1996, 1998, 1999, 2000, 2001a, 2001b, 2002a, 2002b, 2003a, 2003b, 2005), and from 0 to 290°C (Lassin et al., 2015; Christov & Moller, 2004a, 2004b; Moller et al, 2006; Christov, 2007, 2009a, 2009b, 2012).

The thermodynamic properties of the alkali fluoride aqueous solutions are of great interest for the knowledge and better understanding of the physical-chemical processes governing industrial and environmental processes. The study of behavior of solid fluorides and hydrogen fluorides in aqueous solutions play an important role in industrial applications. Thus, the solid potassium fluoride is widely used in the metallurgy, glass fiber optics industry and ceramic manufacturing processes. In the agriculture, KF(s) has been employed in a range of applications, including pesticides, herbicides, and insecticides (El Guendouzi & Faridi, 2020). The solution and solid-liquid equilibrium models for fluoride systems are also of high importance in development of technology for production of fluoride compounds needed in pharmaceutical industry. Models for high soluble solid fluorides (s.a. $\text{KF} \cdot 2\text{H}_2\text{O}(\text{s})$, $\text{RbF}(\text{s})$, and $\text{CsF}(\text{s})$) are of interest in development of strategies and programs for nuclear waste sequestration.

In this study we developed well validated thermodynamic models for solution behavior and solid-liquid equilibrium for all fluoride binary systems, for which activity data are available. The subject of modeling study are 5 fluoride systems of the type 1-1 ($\text{HF-H}_2\text{O}$, $\text{NaF-H}_2\text{O}$, $\text{KF-H}_2\text{O}$, $\text{RbF-H}_2\text{O}$, and $\text{CsF-H}_2\text{O}$) and one of 1-2 type ($\text{H}_2\text{SiF}_6\text{-H}_2\text{O}$) from low to very high concentration ($> 35 \text{ m}$) at 298.15 K. Models are developed on the basis of Pitzer ion interactions approach (Pitzer, 1973, 1991).

Methodology

1. Pitzer's equations

The Pitzer's equations (1 to 4) are described and widely discussed in the literature (Harvie et al., 1984; Moller et al., 2006; Christov and Moller, 2004a, 2004b; Christov, 2005). Here only the expression for the activity coefficient of the interaction of cation (M) with other solutes, $\gamma_{(M+)}$ is given:

$$\ln \gamma_M = z_M^2 F + \sum_a m_a (2B_{Ma}(I) + ZC_{Ma}) + \sum_c m_c (2\Phi_{Mc} + \sum_a m_a \psi_{Mca}) + \sum_a \sum_{a' < a} m_a m_{a'} \psi_{Maa'} + |z_M| \left| \sum_c \sum_a m_c m_a C_{ca} \right. \\ \left. + \sum_n m_n (2\lambda_{nM}) + \sum_n \sum_a m_n m_a \zeta_{naM} \right. \quad (1)$$

Equation (1) is symmetric for anions. The subscripts c and a in eqn. 1 refer to cations and anions, and m is their molality; z is the charge of the M^+ ion. B and Φ represent measurable combinations of the second virial coefficients; C and ψ represent measurable combinations of third virial coefficients. B and C are parameterized from single electrolyte data, and Φ and ψ are parameterized from mixed solution data. The function F is the sum of the Debye-Hueckel term,

$$-A^\phi \left[\sqrt{I} / (1 + b\sqrt{I}) + (2/b) (\ln(1 + b\sqrt{I})) \right], \quad (2)$$

and terms with the derivatives of the second virial coefficients with respect to ionic strength (see Harvie et al., 1984). In Eq. (2), b is a universal empirical constant assigned to be equal to 1.2. A^ϕ (Debye-Hückel limiting law slope for the osmotic coefficient) is a function of temperature, density and the dielectric constant of water (Christov and Moller, 2004b).

For the interaction of any cation M and any anion X in a binary system MX -H₂O, Pitzer assumes that in Eq. (1) B has the ionic strength dependent form:

$$B_{MX} = \beta_{MX}^{(0)} + \beta_{MX}^{(1)} g(\alpha_1 \sqrt{I}) + \quad (3)$$

$$+ \beta_{MX}^{(2)} g(\alpha_2 \sqrt{I}), \quad (3A)$$

where $g(x) = 2[1 - (1 + x)e^{-x}] / x^2$ with $x = \alpha_1 \sqrt{I}$ or $\alpha_2 \sqrt{I}$. α terms are function of electrolyte type and does not vary with concentration or temperature.

In Eq. 1, the Φ terms account for interactions between two ions i and j of like charges. In the expression for Φ ,

$$\Phi_{ij} = \theta_{ij} + {}^E\theta_{ij}(I), \quad (4)$$

θ_{ij} is the only adjustable parameter. The ${}^E\theta_{ij}(I)$ term accounts for electrostatic unsymmetric mixing effects that depend only on the charges of ions i and j and the total ionic strength. The ψ_{ijk} parameters are used for each triple ion interaction where the ions are not all of the same sign. Their inclusion is generally important for describing solubilities in concentrated multicomponent systems. Therefore, according to the basic Pitzer

equations, at constant temperature and pressure, the solution model parameters to be evaluated are: 1) pure electrolyte $\beta^{(0)}$, $\beta^{(1)}$, and C^ϕ for each cation-anion pair; 2) mixing θ for each unlike cation-cation or anion-anion pair; 3) mixing ψ for each triple ion interaction where the ions are all not of the same sign.

2. The $\beta^{(2)}$ parameter (Eqn. 3A) for 2-2 type of electrolytes

Pitzer and Mayorga (1973) did not present analysis for any 2-2 (e.g. $\text{MgSO}_4\text{-H}_2\text{O}$) or higher {e.g. 3-2: $\text{Al}_2(\text{SO}_4)_3\text{-H}_2\text{O}$ } electrolytes. Indeed, they found that three β^0 , β^1 , and C^ϕ parameters approach (see Eqn. 1) could not accurately fit the activity data for these types of solutions. For these electrolytes mean activity ($\gamma_\pm$) and osmotic (ϕ) coefficients drop very sharply in dilute solutions, while showing a very gradual increase, with a very wide minimum at intermediate concentration. Pitzer concluded that this behaviour is due to ion association reactions and that the standard approach with three evaluated solution parameters cannot reproduce this behaviour. This lead to a further (Pitzer & Mayorga, 1974) modification to the original equations for the description of binary solutions: parameter $\beta^2(\text{M},\text{X})$, and an associated $\alpha_2\sqrt{I}$ term are added to the B_{MX} expression (see Eqn. (3A)). Pitzer presented these parameterizations assuming that the form of the functions (i.e. 3 or 4 β and C^ϕ values, as well as the values of the α terms) vary with electrolyte type. For binary electrolyte solutions in which either the cationic or anionic species are univalent (e.g. NaCl , Na_2SO_4 , or MgCl_2), the standard Pitzer approach use 3 parameters (i.e. omit the β^2 term) and α_1 is equal to 2.0. For 2-2 type of electrolytes the model includes the β^2 parameter and α_1 equals to 1.4 and α_2 equals to 12. This approach provides accurate models for many 2-2 binary sulfate (Pitzer & Mayorga, 1974; Christov, 1999, 2000, 2003a) and selenate (Christov, 2003a) electrolytes, giving excellent representation of activity data covering the entire concentration range from low molality up to saturation and beyond.

3. Inclusion of “standard Pitzer approach” $\beta^{(2)}$ parameter into a models for 1-1, 2-1, 3-1, 4-1, 1-2, 1-3, and 3-2 type of electrolytes

To describe the high concentration solution behaviour of systems showing a “smooth” maximum on $\gamma_\pm$ vs. m dependence, and to account strong association reactions at high molality, Christov (1996, 1999, 2000, 2005) used a very simple modelling technology: introducing into a model a fourth ion interaction parameter from basic Pitzer theory $\{\beta^{(2)}\}$, and varying the values of α_1 and α_2 terms. The author also found that by variation of the values of α_1 and α_2 terms it is possible to vary the concentration range of binary solutions at which association reactions become more important and should be account by introducing $\beta^{(2)}$ parameter. According to Christov (2005), model which uses $\alpha_1 = 1.4$ and $\alpha_2 = 12$ accounts association only at low molality solutions (see Christov and Moller (2004b) for $\text{Ca}(\text{OH})_2\text{-H}_2\text{O}$ model). According to previous studies of one of the authors (Christov) an approach with 4 ion interaction parameters ($\beta^{(0)}$, $\beta^{(1)}$, $\beta^{(2)}$, and C^ϕ), and

accepting $\alpha_1 = 2$, and varying in α_2 values can be used for solutions for which ion association occurs in high molality region. This approach was used for binary electrolyte systems of different type: 1-1 type {such as $\text{HNO}_3\text{-H}_2\text{O}$ (Christov, 2005), $\text{LiNO}_3\text{-H}_2\text{O}$ (Donchev and Christov, 2020), and $\text{LiCl-H}_2\text{O}$ (Lassin et al., 2015)}, 2-1 {such as $\text{NiCl}_2\text{-H}_2\text{O}$, $\text{CuCl}_2\text{-H}_2\text{O}$, $\text{MnCl}_2\text{-H}_2\text{O}$, $\text{CoCl}_2\text{-H}_2\text{O}$: (Christov, 1996, 1999, 2000); $\text{Ca(NO}_3)_2\text{-H}_2\text{O}$: (Lach et al., 2018); $\text{UO}_2(\text{NO}_3)_2\text{-H}_2\text{O}$ (Lassin et al., 2020)}, 1-2 {such as $\text{Na}_2\text{Cr}_2\text{O}_7\text{-H}_2\text{O}$: (Christov, 2001a); $\text{K}_2\text{Cr}_2\text{O}_7\text{-H}_2\text{O}$: (Christov, 1998)}, 3-1 {such as $\text{FeCl}_3\text{-H}_2\text{O}$: (Andre et al., 2019); $\text{Ln(NO}_3)_3(\text{aq})$: (Guignot et al., 2019)}, and 3-2 {such as $\text{Al}_2(\text{SO}_4)_3\text{-H}_2\text{O}$, $\text{Cr}_2(\text{SO}_4)_3\text{-H}_2\text{O}$, and $\text{Fe}_2(\text{SO}_4)_3\text{-H}_2\text{O}$: (Christov, 2001b, 2002a, 2002b, 2003b, 2005)}. The resulting models reduce the sigma values of fit of experimental activity data, and extend the application range of models for binary systems to the highest molality, close or equal to molality of saturation {m(sat)}, and in case of data availability: up to supersaturation.

Results and discussions. Model parameterization and validation of models for binary fluoride systems

The activity data for fluoride systems are presented by Mikulin (1968) (for $\text{NaF-H}_2\text{O}$, $\text{KF-H}_2\text{O}$, $\text{RbF-H}_2\text{O}$, and $\text{CsF-H}_2\text{O}$ solutions), by Hammer and Wu (1972) (for all 5 systems under study $\text{HF-H}_2\text{O}$, $\text{NaF-H}_2\text{O}$, $\text{KF-H}_2\text{O}$, $\text{RbF-H}_2\text{O}$, and $\text{CsF-H}_2\text{O}$ solutions), by El Guendouzi and Faridi (2020) (for $\text{KF-H}_2\text{O}$), El Guendouzi et al. (2015) for $\text{H}_2\text{SiF}_6\text{-H}_2\text{O}$, and by Robinson and Stokes (1959) (for $\text{NaF-H}_2\text{O}$, and $\text{KF-H}_2\text{O}$). The data of Mikulin (1968) and Hammer and Wu (1972) are in excellent agreement for $\text{NaF-H}_2\text{O}$, $\text{KF-H}_2\text{O}$, $\text{RbF-H}_2\text{O}$, and $\text{CsF-H}_2\text{O}$ systems. The low molality data of Robinson and Stokes (1959) are also in excellent agreement with the data of Mikulin and Hammer and Wu. Mikulin reported data not only for unsaturated solutions, but also at saturation point. El Guendouzi and Faridi (2020) reported a little higher values for the osmotic coefficients data at high molality in $\text{KF-H}_2\text{O}$ solutions, compare to the data of Mikulin (1968), and Hammer and Wu (1972) (max diff. of 0.04). For $\text{HF-H}_2\text{O}$ solutions the only available mean activity ($\gamma_{\pm}$) and osmotic (ϕ) coefficients data are those of Hammer and Wu (1972). However, our preliminary test shows that there is some inconsistency between the ϕ data at low molality (< 1 m), obtained on the basis of E.M.F. experiments, and the ϕ data at higher molality, obtained using isopiestic experiments. Therefore, the model for $\text{HF-H}_2\text{O}$ solutions have been parameterized on the basis of activity coefficients ($\gamma_{\pm}$) data of Hammer and Wu. For the remaining four 1-1 fluoride systems the osmotic (ϕ) coefficients data for unsaturated, saturated and super-saturated (for $\text{KF-H}_2\text{O}$ system) solutions from Mikulin (1968) are used to construct the models. The osmotic (ϕ) coefficients data up to 3.0 m of El Guendouzi et al. (2015) are used to parameterize the model for $\text{H}_2\text{SiF}_6\text{-H}_2\text{O}$ system up to 10 m.

In parameterization process we used the value of Debye-Hückel term (A^{ϕ}) equals to 0.39147 at 25°C , as given in Christov (2009a, 2009b, 2012). To construct the models we used different versions of standard molality-based Pitzer approach. It was established that for 4 of systems under study ($\text{NaF-H}_2\text{O}$, $\text{KF-H}_2\text{O}$,

RbF-H₂O, and CsF-H₂O) application of extended approach with 4 parameters (β^0 , β^1 , β^2 and C^ϕ) and variation of α_2 term in fundamental Pitzer equations leads to the lowest values of standard model-experiment deviation (see paragraph 3 in the Methodology). To account association effects at low and high concentrated HF-H₂O solutions the model for this system has been constructed by variation of both (α_2 and α_2) terms.

On next figures (A, B, C, D, E, F, and G) we compare model calculated (lines) mean activity coefficients (Fig. A) and osmotic coefficients (ϕ) (Fig. B, C, D, E, F, G) of HF (Fig. A), of H₂SiF₆ (Fig. B), NaF (Fig. C), KF (Fig. D), RbF (Fig. E) and CsF (Fig. F and G) in binary solutions HF-H₂O, H₂SiF₆-H₂O, NaF-H₂O, KF-H₂O, RbF-H₂O, and CsF-H₂O) against molality at T = 298.15 K with osmotic (ϕ) coefficients data, and activity coefficients ($\gamma_{\pm}$) recommendations in literature (symbols). The (E) figure is an enlargement of the low molality corner for CsF-H₂O. Heavy solid lines represent the predictions of the developed in this study and accepted models. Dashed-dotted, and dashed lines represent the predictions of the reference models of Kim and Frederick (1988) (Fig. A), of El Guendouzi et al. (2015) (Fig. B), of Pitzer and Mayorga (1973) (Fig. C, D, E, F, and G), and of El Guendouzi and Faridi (2020) (Fig. D). The experimental data are taken from Mikulin (1968) (open triangles on all figures and open squares on Fig. F and G), Hammer and Wu (1972) (open squares on Fig. B, C, D, and F), El Guendouzi and Faridi (2020) (crosses on Fig. E), and El Guendouzi et al. (2005) (open diamonds on Fig. B). The vertical lines on figures denote the molality of stable crystallization of solid fluoride phases (from Mikulin, 1968).

As shown on figures the predictions of new developed here models are in excellent agreement with experimental osmotic coefficients data, and with recommendations on activity coefficients in binary solutions from low (see Fig. F) to very high concentration: up to 20 mol.kg⁻¹ in HF-H₂O, up to 17.5 mol.kg⁻¹ in KF-H₂O, up to 22.4 mol.kg⁻¹ in RbF-H₂O and up to 35.6 mol. kg⁻¹ in CsF-H₂O. With only one exception (for NaF-H₂O), the models developed here are in a pure agreement with models reported in literature. For NaF-H₂O system our model is in very good agreement with the low molality model of Pitzer and Mayorga (1973), and the data. The model of Kim & Frederick (1988) predicts much higher activity coefficients for whole concentration range (up to 20 m) for HF-H₂O solutions, compare to the data and our new model. For H₂SiF₆-H₂O system our model predicts a little higher osmotic coefficients values at high molality compare to the model of El Guendouzi et al. (2005). New data are needed to prove the correctness of model predictions.

A)

B)

C)

D)

E)

F)

G)

Fig. 1. (A, B, C, D, E, F, G) A comparison of model calculated (lines) mean activity coefficients (Fig. A) and osmotic coefficients (ϕ) (Fig. B, C, D, E, F, G) of HF (Fig. A), H_2SiF_6 (Fig. B), NaF (Fig. C), KF (Fig. D), RbF (Fig. E) and CsF (Fig. F and G) in binary solutions HF- H_2O , H_2SiF_6 - H_2O , NaF- H_2O , KF- H_2O , RbF- H_2O , and CsF- H_2O against molality at $T = 298.15$ K with osmotic (ϕ) coefficients data, and activity coefficients ($\gamma_{\pm}$) recommendations in literature (symbols). The (G) figure is an enlargement of the low molality corner. Heavy solid lines represent the predictions of the developed in this study and accepted models. Dashed-dotted, and dashed lines represent the predictions of the reference models of Kim and Frederick (1988) (Fig. A), of El Guendouzi et al. (2015) (Fig. B), of Pitzer and Mayorga (1973) (Fig. C, D, E, F and G), and of El Guendouzi and Faridi (2020) (Fig. D). The experimental data are taken from Mikulin (1968) (open triangles on all figures and open squares on Fig. F and G), Hammer and Wu (1972) (open squares on Fig. B, C, D and F), El Guendouzi and Faridi (2020) (crosses on Fig. E), and El Guendouzi et al. (2015) (open diamonds on Fig. B). The vertical lines on figures denote the molality of stable crystallization of solid fluoride phases (from Mikulin, 1968)

For KF- H_2O solutions the predictions of the models of Pitzer and Mayorga (1973) and of El Guendouzi and Faridi (2020) are in good agreement with data and our model only up to ≈ 10 m. At higher molality these models predict higher values of osmotic coefficients. The comparison on Fig. (E) and (F) shows that for RbF- H_2O and CsF- H_2O systems the model of Pitzer and Mayorga (1973) reproduced well the data only at low molality (up to ≈ 3.0 m for RbF- H_2O , and up to ≈ 7 m for CsF- H_2O). Therefore, it can be

concluded that with only two exceptions (the model of Pitzer and Mayorga (1973) for NaF-H₂O, and the model of El Guendouzi et al. (2005) for H₂SiF₆-H₂O) all other models given in literature are concentration restricted and can't be used for solubility calculations and correct determination of properties of crystallizing solid phases.

On the basis of evaluated in this study binary parameters ($\beta^{(0)}$, $\beta^{(1)}$, $\beta^{(2)}$, and C^ϕ) we also determine water activity (a_w) and Deliquescence Relative Humidity (DRH (%)) (see eqn. 5) of solid phases crystallizing from saturated binary fluoride solutions [NaF(s), KF.2H₂O(s), RbF(s), and CsF(s)].

Within the solid-liquid equilibrium model, relative humidity is related to water activity, a_w (Clegg et al., 1998; Christov, 2009a,b; 2012):

$$a_w = P_w / P_w^0 = \text{RH}/100, \quad (5)$$

where P_w and P_w^0 are the vapor pressure of the saturation solution and pure water, respectively, at given temperature. The results of calculations are given in Table 1. The model DRH predictions are in excellent agreement with the experimental data determined using isopiestic method, and given in Mikulin (1968). According to model calculations the solid-liquid phase change of CsF(s) occurs at lowest relative humidity (RH) of environment ($\approx 2.5\%$).

As a next step, using the accepted new developed parameterizations, and experimentally determined molalities ($m(\text{sat})$) of the saturated binary solutions (Mikulin, 1968) we calculate the logarithm of the thermodynamic solubility product ($\ln K_{\text{sp}}^0$) of 4 solid phases crystallizing from saturated binary fluoride solutions at $T = 298.15 \text{ K}$ (eqn. (6)).

$$\begin{aligned} K_{\text{sp}}^0(\text{NaF}) &= \gamma_{(\pm)}(\text{sat})^2 \cdot m(\text{sat})^2 \\ K_{\text{sp}}^0(\text{KF} \cdot 2\text{H}_2\text{O}) &= \gamma_{(\pm)}(\text{sat})^2 \cdot m(\text{sat})^2 \cdot a_w(\text{sat})^2 \end{aligned} \quad (6)$$

The model calculations are given in Table 1. The small $\ln K_{\text{sp}}^0$ difference (of 0.26 logarithm units) for KF.2H₂O (s) compare to calculations of El Guendouzi and Faridi (2020) is mainly due on the different $m(\text{sat})$ values used in calculations (see eqn. (6) and Table 1). Note that the widely used databases of Pitzer (1991), Pitzer and Mayorga (1973, 1974), and Kim and Frederick (1988) do not consider solid phases.

Table 1. Comparison between model calculated and recommended values of thermodynamic properties for fluoride solid phases crystallizing from saturated binary solutions at $T = 298.15$ K: 1) Deliquescence Relative Humidity [DRH (%) = $a_w^s \times 100$; where a_w^s is activity of water at saturation], and 2) logarithm of the thermodynamic solubility product, K_{sp}^o ,

Salt composition	m(sat) (exp) ^a (mol.kg ⁻¹)	DRH (%)		lnK _{sp} ^o	
		Calculated	Ref. data ^a	Calculated	El Guendouzi and Faridi ^b
NaF(cr)	1.0	96.91	96.90	-1.115	
KF.2H ₂ O(cr)	16.68	31.33	31.10	5.84	6.1051 (m(sat)=17.5)
RbF(cr)	22.4	15.20	14.50	10.801	
CsF(cr)	35.6	2.46	4.00	14.74	

^afrom Mikulin (1968)

^bfrom El Guendouzi and Faridi (2020)

Acknowledgement

The work was supported by the European Regional Development Fund within the Operational Programme “Science and Education for Smart Growth 2014-2020” under the Project CoE “Universities for Science, Informatics and Technologies in e-Society (UNITE) BG05M2OP001-1.001-0004”, as well as by the Shumen University Research Fund (project RD 08-131/04.02.2021).

References

- [1] André L., Christov C., Lassin A., Azaroual M. A thermodynamic model for solution behavior and solid-liquid equilibrium in Na-K-Mg-Ca-Al(III)-Fe(III)-Cr(III)-Cl-H₂O system from low to very high concentration at 25°C, *Acta Scientifica Naturalis*, **2019**, 6(1), 26-36.
- [2] Christov C. Thermodynamic study of the co-crystallization of 2RbCl.NiCl₂.2H₂O and 2RbCl.MnCl₂.2H₂O at the temperature 298.15 K, *J. Chem. Thermodynamics*, **1996**, 28, 743-752.
- [3] Christov C. Thermodynamic study of the KCl-K₂SO₄-K₂Cr₂O₇-H₂O system at the temperature 298.15 K, *CALPHAD*, **1998**, 22, 449-457.
- [4] Christov C. Study of (m₁KCl + m₂MeCl₂)(aq), and (m₁K₂SO₄ + m₂MeSO₄)(aq) where m denotes molality

- and Me denotes Cu or Ni, at the temperature 298.15 K, *J. Chem. Thermodynamics*, **1999**, 31, 71-83.
- [5] Christov C. Thermodynamic study of the Na-Cu-Cl-SO₄-H₂O system at the temperature 298.15 K, *J. Chem. Thermodynamics*, **2000**, 32, 285-295.
- [6] Christov C. Thermodynamic study of the NaCl-Na₂SO₄-Na₂Cr₂O₇-H₂O system at the temperature 298.15 K, *CALPHAD*, **2001a**, 25, 11-17.
- [7] Christov C. Thermodynamic study of the K-Mg-Al-Cl-SO₄-H₂O system at the temperature 298.15 K, *CALPHAD*, **2001b**, 25, 445-454.
- [8] Christov C. Thermodynamics of formation of ammonium, sodium, and potassium alums and chromium alums, *CALPHAD*, **2002a**, 26, 85-94.
- [9] Christov C. Thermodynamic study of quaternary systems with participation of ammonium and sodium alums and chromium alums, *CALPHAD*, **2002b**, 26, 341-352.
- [10] Christov C. Thermodynamics of formation of double salts M₂SO₄.MeSO₄.6H₂O and M₂SeO₄.MeSeO₄.6H₂O where M denotes Rb, or Cs, and Me denotes Co, Ni or Zn, *J. Chem. Thermodynamics*, **2003a**, 35, 1775-1792.
- [11] Christov C. Thermodynamic study of the co-crystallization of ammonium, sodium and potassium alums and chromium alums, *CALPHAD*, **2003b**, 27, 153-160.
- [12] Christov C. Thermodynamics of formation of double salts and solid solutions from aqueous solutions, *J. Chem. Thermodynamics*, **2005**, 37, 1036-1060.
- [13] Christov C. An isopiestic study of aqueous NaBr and KBr at 50°C. Chemical Equilibrium model of solution behavior and solubility in the NaBr-H₂O, KBr-H₂O and Na-K-Br-H₂O systems to high concentration and temperature, *Geochim. Cosmochim. Acta*, **2007**, 71, 3357-3369.
- [14] Christov C. Isopiestic Determination of the osmotic coefficients of aqueous MgCl₂-CaCl₂ Mixed solution at 25°C and 50°C. Chemical equilibrium model of solution behavior and solubility in the MgCl₂-H₂O, and MgCl₂-CaCl₂-H₂O systems to high concentration at 25°C and 50°C, *J. Chem. Eng. Data*, **2009a**, 54, 627-635.
- [15] Christov C. Chemical equilibrium model of solution behavior and solubility in the MgCl₂-H₂O, and HCl-MgCl₂-H₂O systems to high concentration from 0°C to 100°C, *J. Chem. Eng. Data*, **2009b**, 54, 2599-2608.
- [16] Christov C. Study of bromide salts solubility in the (m₁KBr + m₂CaBr₂)(aq) system at T = 323.15 K. Thermodynamic model of solution behavior and solid-liquid equilibria in the ternary (m₁KBr + m₂CaBr₂)(aq), and (m₁MgBr₂ + m₂CaBr₂)(aq), and in quinary {Na+K+Mg+Ca+Br+H₂O} systems to high concentration and temperature, *J. Chem. Thermodynamics*, **2012**, 55, 7-22.
- [17] Christov C., Moller N. Chemical equilibrium model of solution behavior and solubility in the H-Na-K-

- Cl-OH-HSO₄-SO₄-H₂O system to high concentration and temperature, *Geochim.Cosmochim. Acta*, **2004a**, 68, 1309-1331.
- [18] Christov C., Moller N. A chemical equilibrium model of solution behavior and solubility in the H-Na-K-Ca-Cl-OH-HSO₄-SO₄-H₂O system to high concentration and and temperature, *Geochim.Cosmochim. Acta*, **2004b**, 68, 3717-3739.
- [19] Clegg S., Brimblecombe P., Wexler A. Thermodynamic model of the system H⁺-NH₄⁺- Na⁺- SO₄²⁻-NO₃⁻-Cl⁻-H₂O at 298.15 K. *J. Phys. Chem.*, **1998**, A102, 2155-2171.
- [20] Donchev S., Christov C. Development of Accurate Chemical Thermodynamic Database for Geochemical Storage of Nuclear Waste. Part I: Models for Predicting Solution Properties and Solid-Liquid Equilibrium in Binary Nitrate Systems of the Type 1-1, *Ecologia Balkanica*, Special Edition 3, **2020**, pp. 195-210.
- [21] El Guendouzi M., Faridi, Thermodynamic properties and solubility of potassium fluoride in aqueous solutions at various temperatures, *J. Fluorine Chem.*, **2020**, 235, 109558 (<https://doi.org/10.1016/j.jfluchem.2020.109558>).
- [22] El Guendouzi M., Rifai A., Skafi M. Properties of fluoride in wet phosphoric acid processes: Fluorosilicic acid in an aqueous solution of H₂SiF₆-H₂O at temperatures ranging from 298.15 K to 353.15 K, *Fluid Phase Equilibria*, **2015**, 396, 43-49.
- [23] Guignot S., Lassin A., Christov C., Lach A., André L., Henocq P. Modelling the osmotic and activity coefficients of lanthanide nitrate aqueous solutions at 298.15 K from low molalities to (super) saturation, *J. Chem. Eng. Data*, **2019**, 64(1), 345-359.
- [24] Hamer W.J., Wu Y-C. Osmotic coefficients and mean activity coefficients of uni-univalent electrolytes in water at 25°C. *J. Phys. Chem. Ref. Data*, **1972**, 1, 1047-1099.
- [25] Harvie C., Moller N., Weare J. The prediction of mineral solubilities in natural waters: The Na-K-Mg-Ca-H-Cl-SO₄-OH-HCO₃-CO₃-CO₂-H₂O system from zero to high concentration at 25°C. *Geochim. Cosmochim. Acta*, **1984**, 48, 723-751.
- [26] Kim H.-T., Frederick W. Evaluation of Pitzer ion interaction parameters of aqueous electrolytes at 25°C. 1. Single salt parameters. *J. Chem. Eng. Data*, **1988**, 33, 177-184.
- [27] Kolev H., Tyuliev G., Christov C., Kostov K. Experimental study of the surface chemical composition of sea salt crystallized at evaporation of sea water under natural conditions, *Bulgarian Chem. Commun.* **2013**, 45, 584-591.
- [28] Lach A., André L., Guignot S., Christov C., Henocq P., Lassin A. A Pitzer parameterization to predict solution properties and salt solubility in the H-Na-K-Ca-Mg-NO₃-H₂O system at 298.15 K, *J. Chem. Eng. Data*, **2018**, 63, 787-800.

- [29] Lassin A., Christov C., André L., Azaroual M. A thermodynamic model of aqueous electrolyte solution behavior and solid-liquid equilibrium in the Li-H-Na-K-Cl-OH-H₂O system to very high concentrations (40 Molal) and from 0 to 250°C. *Amer. J. Sci.*, **2015**, 315, 204–256.
- [30] Lassin, A., Guignot, S., Lach, A., Christov, C., André, L., Madé, B. Modeling the solution properties and mineral-solution equilibria in radionuclide-bearing aqueous nitrate systems. Application to binary and ternary systems containing U, Th or lanthanides, at 25°C, *J. Chem. Eng. Data*, **2020**, 65, 3613-3626.
- [31] Mikulin G. *Voprosy Fizicheskoi Khimii Electrolytov*, Izd. Khimiya, St. Petersburg, **1968**, 417 p.
- [32] Moller N., Christov C., Weare J. Thermodynamic models of aluminumsilicate mineral solubility for application to enhanced geothermal systems. *Proceedings from 31th Workshop on Geothermal Reservoir Engineering*, Stanford University, Stanford, California, January 30 - February 1, **2006** (8 pages).
- [33] Park J.-H., Christov C., Ivanov A., Molina M. On OH uptake by sea salt under humid conditions, *Geophysical Research Letters*, **2009**, 36 LO2802, DOI:10.1029/2009GL036160.
- [34] Pitzer K. S. Thermodynamics of Electrolytes. I. Theoretical Basis and General Equations, *J. Phys. Chem.*, **1973**, 77(2), 268-277.
- [35] Pitzer K.S. Ion Interaction Approach: Theory and Data Correlation. Chapter 3 of *Activity Coefficients in Electrolyte Solutions*. 2nd Edition. Pitzer, K.S., ed. Boca Raton, Florida: CRC Press. TIC: 251799. 123206 , **1991**.
- [36] Pitzer K.S., Mayorga G. Thermodynamics of electrolytes. II. Activity and osmotic coefficients for strong electrolytes with one or both ions univalent. *J. Phys. Chem.*, **1973**, 77, 2300-2308.
- [37] Pitzer, K.S., Mayorga G. Thermodynamics of electrolytes. III. Activity and osmotic coefficients for 2-2 electrolytes. *J. Soln. Chem.*, **1974**, 3, 539-546.
- [38] Robinson R. and Stokes R. *Electrolyte Solutions*, 2nd ed.; Butterworths, London, **1959**.