Comments on the Half Logistic Inverse Rayleigh (HLIR) cdf with “Polynomial Variable Transfer”. Some Applications

Nikolay Kyurkchiev, Anton Iliev, Asen Rahnev

Faculty of Mathematics and Informatics, University of Plovdiv Paisii Hilendarski, 24 Tzar Asen Str., 4000 Plovdiv, Bulgaria

E-mails: nkyurk@uni-plovdiv.bg aii@uni-plovdiv.bg assen@uni-plovdiv.bg

Abstract: *On the base of the Half Logistic – G family of distributions proposed by Cordeiro, Alizadeh and Marinho [2] some mathematical properties are investigated by Almarashi et al. [1]. We study the “saturation” to the horizontal asymptote: $t=1$ by the new growth function $M(t)$ in the Hausdorff sense. Similar to our previous studies [3-6], in this article we will define and analyze in detail the new family. We will call this family the “Half-Logistic-Inverse-Rayleigh cdf with Polynomial Variable Transfer” (HLIRPVT) cdf. Section 3 shows the potentiality of proposed new model under four real data sets. Some numerical examples using CAS MATHEMATICA are given.*

Keywords: *Half-Logistic-Inverse-Rayleigh (HLIR) cdf, Half-Logistic-Inverse-Rayleigh cdf with Polynomial Variable Transfer (HLIRPVT) cdf, “Saturation” in Hausdorff sense, Hausdorff distance, Upper and lower bounds.*

1. Introduction and preliminaries

Definition 1. In [1] authors proposed the following new Half-Logistic-Inverse-Rayleigh (HLIR) cdf:

$$(1) \quad M(t) = \frac{1 - \left(1 - e^{-\left(\frac{\alpha}{t}\right)^2}\right)^\lambda}{1 + \left(1 - e^{-\left(\frac{\alpha}{t}\right)^2}\right)^\lambda},$$

where $t > 0, \alpha > 0, \lambda > 0$, and proved the following inequalities for any, $t, \alpha, \lambda > 0$:

$$(2) \quad M_{**}(t) < M(t) < M_*(t),$$

$$M_*(t) = 1 - \left(1 - e^{-\left(\frac{\alpha}{t}\right)^2} \right)^\lambda ; \quad M_{**}(t) = \frac{t^{2\lambda}}{t^{2\lambda} + \alpha^{2\lambda}} M_*(t),$$

where $M_*(t)$ is the Exponentiated Inverted Rayleigh cdf, and $M_{**}(t)$ is a weighted version of it, which remains a valid cdf.

During the last two decades or so, various approaches of determining new families of distributions have been introduced for increasing chances of modeling practical data that come from a wide variety of disciplines.

The reader can find a detailed bibliography in article [7].

In this paper we study some properties of the new family.

When studying the intrinsic properties of the $M(t)$, it is also appropriate to study the “saturation” to the horizontal asymptote.

Definition 2. Similar to our previous studies [3-6], in this article we will define and analyze in detail the following new family:

$$(3) \quad M_1(t; a_1, \dots, a_n) = \frac{1 - \left(1 - e^{-\left(\frac{\alpha}{F(t)}\right)^2} \right)^\lambda}{1 + \left(1 - e^{-\left(\frac{\alpha}{F(t)}\right)^2} \right)^\lambda},$$

$$F(t) = \sum_{i=1}^n a_i t^i.$$

We will call this family the “Half-Logistic-Inverse-Rayleigh cdf with Polynomial Variable Transfer” (HLIRPVT) cdf.

The rest of the paper is structured as follows. In Section 2 we study the “saturation” to the horizontal asymptote: $t = 1$ by the growth functions (1) and (3) in the Hausdorff sense [19].

Section 3 shows the potentiality of proposed new model based on the insertion of “correcting amendments” of polynomial-type (3) for the estimation procedure under two real data sets.

2. Main results

1. For the “saturation” d_1 to the horizontal asymptote: $t = 1$ by the growth function (1) in the Hausdorff sense we have

$$(4) \quad M(d_1) = \frac{1 - \left(1 - e^{-\left(\frac{\alpha}{d_1}\right)^2}\right)^\lambda}{1 + \left(1 - e^{-\left(\frac{\alpha}{d_1}\right)^2}\right)^\lambda} = 1 - d_1,$$

or, equivalently,

$$(5) \quad G(d_1) := \frac{1}{\left(1 - e^{-\left(\frac{\alpha}{d_1}\right)^2}\right)^\lambda} + 1 - \frac{2}{d_1} = 0.$$

The following theorem gives upper bound for d_1 .

Theorem 1. The “saturation” d_1 satisfies the following inequality for $\alpha > 0, \lambda > 0$:

$$(6) \quad d_1 < \left(2\alpha^{2\lambda}\right)^{\frac{1}{1+2\lambda}} := d_{1,r}.$$

Proof: We consider the interval $[0, +\infty)$.

Clearly, $G' > 0$ and $G(t)$ is increasing function of $t \in [0, +\infty)$ (Fig. 1).

Fig. 1. The function G for $\lambda = 2.3$ and $\alpha = 0.05$; the “saturation” $d_1 = 0.090098$

Hence, if (5) has a root, then it is unique.

Using the inequalities

$$1 \leq 1 + \left(1 - e^{-\left(\frac{\alpha}{t}\right)^2}\right)^\lambda < 1 + \left(\frac{\alpha}{t}\right)^{2\lambda}; \quad e^\beta \geq 1 + \beta, \quad \beta \in \mathbb{R},$$

we have

$$G(d_1) > \left(\frac{d_1}{\alpha}\right)^{2\lambda} - \frac{2}{d_1} = \frac{d_1^{2\lambda+1} - 2\alpha^{2\lambda}}{\alpha^{2\lambda} d_1}.$$

The positive root of the expression in the numerator is

$$\left(2\alpha^{2\lambda}\right)^{\frac{1}{1+2\lambda}} := d_{1,r},$$

Then $G(d_{1,r}) > 0$.

This completes the proof of the theorem.

2. The one-sided Hausdorff distance d between the Heaviside function $h_{t_0}(t)$ and the model (1) satisfies the relation

$$(7) \quad M(t_0 + d) = 1 - d,$$

where t_0 is the “median level”, i.e., $M(t_0) = \frac{1}{2}$. Some computational examples using model (1) are presented in Figs 2 and 3.

Fig. 2. The model (1) and two-sided estimations (2) for fixed $\alpha = 0.05$, $\lambda = 2.3$, and $t_0 = 0.0508144$,
Housdorf distance $d = 0.0526852$

Fig. 3. The model (1) and two-sided estimations (2) for fixed $\alpha = 0.01$, $\lambda = 0.9$, and $t_0 = 0.0169128$,
Housdorf distance $d = 0.0551343$

From Figs 2, 3 it can be seen that the characteristic d and estimations (2) can be used as “confidence bounds”, which are extremely useful for the specialists in the choice of model for cumulative data approximating in areas of Biostatistics, Population dynamics, Growth theory, Debugging and Test theory, Computer viruses propagation, Insurance mathematics and Financial Mathematics.

3. Investigations on the “Half-Logistic-Inverse-Rayleigh cdf with Polynomial Variable Transfer” (HLIRPVT) cdf (3).

Let t_0 is the “median level”. The one-sided Hausdorff distance d_2 between Heaviside function h_{t_0} and cumulative function (3) satisfies

$$M_1(d_2; a_1, \dots, a_n) = 1 - d_2.$$

The question of finding precise two-sided estimates for the magnitude of the Hausdorff approximation [19] of the Heaviside function with classes of the indicated family $M_1(t; a_1, \dots, a_n)$ remains open.

The task is greatly complicated by the intrinsic properties of the generated class of functions, as well as by the type and location of zeros of polynomial $F(t)$.

Methods for simultaneous approximation of all roots of a given polynomial of degree n can be found in [20, 21].

In some of cases when only a part of all zeros of polynomial are necessary to be known methods given in [22] can be used.

Below we look at some comparisons (Fig. 4) between the Half-Logistic-Inverse-Rayleigh (HLIR) cdf and the “Half-Logistic-Inverse-Rayleigh with Polynomial Variable Transfer” (HLIRPVT) cdf, as well as the ability to approximate specific dynamics data.

Fig. 4. Comparison between the model (1) (dashed) and the new model (3) (red) at fixed $\alpha = 0.001$, $\lambda = 0.95$, $n = 3$, $a_1 = -1.3$, $a_2 = 0.03$, $a_3 = -0.00001$. For the model (3) $t_0 = 0.0125191$, Hausdorff distance $d_2 = 0.0439798$

We conclude that the proposed model $M_1(t; a_1, \dots, a_n)$ has some free parameters leading to greater flexibility in modeling various data types.

3. Some applications

Example 1. The model explored here can be successfully used for approximating propagation data of computer viruses [10-18, 25].

As an example we will approximate Stuxnet spreading. Stuxnet is world's first cyberweapon. On 20 July 2010 Symantec initiate to monitor traffic to the Stuxnet Command and Control (C&C) servers [9]. The worm infected over 200,000 computers.

For the cumulative data from Fig. 5 the fitted model

$$M_1^*(t; a_1, \dots, a_n) = AM_1(t; a_1, \dots, a_n) \text{ for } \lambda = 0.41, \quad \alpha = 0.00157998, \quad n = 5, \\ A = 209000, \quad a_1 = -0.00589269, \quad a_2 = 0.00630298, \quad a_3 = -0.00371303, \\ a_4 = 0.000861294, \quad a_5 = -0.0000764171$$

is presented on Fig. 6.

Fig. 5. Stuxnet propagation from 20 July 2010 to 21 September 2010 [9]

Fig. 6. The fitted model $M_1^*(t; a_1, \dots, a_n) = AM_1(t; a_1, \dots, a_n)$ for the cumulative data from Fig. 5

Example 2. Approximation of the data “Data Zika Virus Infection” [44, 45].

The number of these M&CD diagnoses to be confirmed with a Zika virus infection also grows (green dots; left-hand axis) by 1 new detection (green bars; right-hand axis) to 208 this EW.

Those confirmed Zika virus infections represent 15% of all confirmed M&CD diagnoses and 3% of all suspect diagnoses.

For the cumulative data (Fig. 7) the fitted model

$$M_1^*(t; a_1, \dots, a_n) = AM_1(t; a_1, \dots, a_n)$$

for $n = 3$, $\lambda = 0.91$, $\alpha = 8318.88$, $A = 220$,
 $a_1 = 3134.48$, $a_2 = -416.973$, $a_3 = 25.9594$

is depicted in Fig. 8.

Fig. 7. Laboratory confirmed ZIKV in microcephaly diagnoses [44, 45]

Fig. 8. The fitted model $M_1^*(t; a_1, \dots, a_n) = AM_1(t; a_1, \dots, a_n)$ for the data “Data Zika Virus Infection” [44, 45]

From the attached graph, it can be seen that at the degree of the polynomial $n = 3$, some specific distortions typical of the model analyzed are very well approximated.

Example 3. Approximation of the data "Influenza A(H1N1) (also known as Swine influenza)" [46].

In April 2009 scientific laboratories discover an influenza virus known as H1N1. This virus seems like a combination of viruses from birds, humans and pigs. Later, in 2010 The World Health Organization (WHO) reported the infection caused by H1N1 as a global pandemic. In August 2010 the WHO reported the finishing of pandemic. Since that scientists changed the name of infection and H1N1 virus is now H1N1v.

Fig. 9. Influenza A(H1N1) cases [46]

For the cumulative data from Fig. 9 the fitted model

$$M_1^*(t; a_1, \dots, a_n) = AM_1(t; a_1, \dots, a_n) \text{ for } \lambda = 1.5, \alpha = 21021.1, n = 6, A = 13398, \\ a_1 = 2777.7, a_2 = -69.8471, a_3 = -26.6675, a_4 = 3.21043, a_5 = -0.131274$$

is presented on Fig. 10.

A detailed discussion of the topic the reader can find in the monographic study [5]. For other results, see [23-41, 42, 43].

Fig. 10. The fitted model $M_1^*(t; a_1, \dots, a_n) = AM_1(t; a_1, \dots, a_n)$ for the data "Influenza A(H1N1)" (also known as "Swine influenza") [46]

Example 4. Approximation of the data “Data South Korea: Total Cases COVID-19” from 15.02.2020 to 26.04.2020 [8].

For the cumulative data from Fig. 11 the fitted model $M_1^*(t; a_1, \dots, a_n)$ for

$$\lambda = 1.7109099, \alpha = 1039.51, n = 5, A = 10728, a_1 = 17.2212,$$

$$a_2 = 6.32739, a_3 = -0.267302, a_4 = 0.00374985, a_5 = -0.0000123795$$

is presented on Fig. 12.

Total Coronavirus Cases in South Korea

Fig. 11. “Data South Korea: Total Cases COVID-19” from 15 February 2020 to 26 April 2020 [8]

Fig. 12. The fitted model $M_1^*(t; a_1, \dots, a_n) = AM_1(t; a_1, \dots, a_n)$

for the “Data South Korea: Total Cases COVID-19” from 15 February 2020 to 26 April 2020 [8]

Acknowledgments: Authors would like to thanks to anonymous reviewers for helpful remarks. This work has been accomplished with the financial support by the Grant No BG05M2OP001-1.001-0003, financed by the Science and Education for Smart Growth Operational Program (2014-2020) and co-financed by the European Union through the European structural and Investment funds.

References

1. Almarashi, A., M. Badr, M. Elgarhy, F. Jamal, C. Chesneau. Statistical Inference of the Half-Logistic-Inverse-Rayleigh Distribution. – *Entropy*, Vol. **22**, 2020.
2. Cordeiro, G. M., M. Alizadeh, E. P. R. D. Marinho. The Type I Half-Logistic Family of Distributions. – *J. Stat. Comput. Simul.*, Vol. **86**, 2016, pp. 707-728.
3. Kyurkchiev, N., A. Iliev, A. Rahnev. On the Verhulst Growth Model with “Polynomial Variable Transfer”. Some Applications. – *International Journal of Differential Equations and Applications*, Vol. **19**, 2020, No 1, pp. 15-32.
4. Kyurkchiev, N., A. Iliev, A. Rahnev. On the Half-Logistic Model with “Polynomial Variable Transfer”. Application to Approximate the Specific “Data CORONA VIRUS”. – *International Journal of Differential Equations and Applications*, Vol. **19**, 2020, No 1, pp. 45-61.
5. Kyurkchiev, N. Selected Topics in Mathematical Modeling: Some New Trends (Dedicated to Academician Blagovest Sendov (1932-2020)). LAP LAMBERT Academic Publishing, 2020. ISBN: 978-620-2-51403-3.
6. Kyurkchiev, N., A. Iliev, A. Rahnev, T. Terzieva. On the Extended Half-Logistic Model by H. Bakouch with “Polynomial Variable Transfer”. Application to Approximate the Specific “Data BG COVID-19”, 2020 (to Appear).
7. Bakouch, H. A Family of Extended Half-Distributions: Theory and Applications. *Filomat*, 2020.
8. <https://www.worldometers.info/coronavirus/country/south-korea/>
9. Falliere, N., L. O’Murchu, E. Chien. W32.Stuxnet Dossier. Version 1.4 (February 2011). Symantec Security Response, 2011. 69 p.
10. Kyurkchiev, N., A. Iliev, A. Rahnev, T. Terzieva. A New Analysis of Code Red and Witty Worms Behavior. – *Communications in Applied Analysis*, Vol. **23**, 2019, No 2, pp. 267-285.
11. Iliev, A., N. Kyurkchiev, A. Rahnev, T. Terzieva. Some New Approaches for Modelling Large-Scale Worm Spreading on the Internet. II. – *Neural, Parallel, and Scientific Computations*, Vol. **27**, 2019, No 1, pp. 23-34.
12. Kyurkchiev, N., A. Iliev, A. Rahnev, T. Terzieva. A New Analysis of Cryptolocker Ransomware and Welchia Worm Propagation Behavior. Some Applications. III. – *Communications in Applied Analysis*, Vol. **23**, 2019, No 2, pp. 359-382.
13. Terzieva, T., A. Iliev, A. Rahnev, N. Kyurkchiev. The Lomax-D-Generalized-Weibull Cumulative Sigmoid with Applications to the Theory of Computer Viruses Propagation. IV. – *Neural, Parallel, and Scientific Computations*, Vol. **27**, 2019, No 3, 4, pp. 141-150.
14. Terzieva, T., A. Iliev, A. Rahnev, N. Kyurkchiev. On a Powerful Transmuted Odd Log-Logistic-Gumbell Model with Applications to the Theory of Computer Viruses Propagation. V. – *Communications in Applied Analysis*, Vol. **23**, 2019, No 3, pp. 441-451.
15. Terzieva, T., A. Iliev, A. Rahnev, N. Kyurkchiev. Comments on a New Hyperbolic Sine-Weibull Model with Applications to the Theory of Computer Viruses Propagation. VI. – *International Journal of Differential Equations and Applications*, Vol. **18**, 2019, No 1, pp. 137-146.
16. Malinova, A., O. Rahneva, A. Golev, V. Kyurkchiev. Investigations on the Odd-Burr-III-Weibull Cumulative Sigmoid. Some Applications. – *Neural, Parallel, and Scientific Computations*, Vol. **27**, 2019, No 1, pp. 35-45.
17. Rahneva, O., T. Terzieva, A. Golev. Investigations on the Zubair-Family with Baseline Ghosh-Bourguignon’s Extended Burr XII Cumulative Sigmoid. Some Applications. – *Neural, Parallel, and Scientific Computations*, Vol. **27**, 2019, No 1, pp. 11-22.

18. Angelova, E., A. Golev, T. Terzieva, O. Rahneva. A Study on a Hyper-Power-Logistic Model. Some Applications. – Neural, Parallel and Scientific Computations, Vol. **27**, 2019, No 1, pp. 45-57.
19. Sendov, B. Hausdorff Approximations. Boston, Kluwer, 1990.
20. Sendov, B., A. Andreev, N. Kyurkchiev. Numerical Solution of Polynomial Equations. – In: Handbook of Numerical Analysis. III. P. Ciarlet, J. Lions, Eds. Amsterdam, Elsevier Science Publ., 1994.
21. Kyurkchiev, N. Initial Approximation and Root Finding Methods. – In: WILEY-VCH Verlag Berlin GmbH. Vol. **104**. 1998.
22. Iliev, A., N. Kyurkchiev. Nontrivial Methods in Numerical Analysis: Selected Topics in Numerical Analysis. Saarbrücken, LAP Lambert Academic Publishing, 2010. ISBN: 978-3-8433-6793-6.
23. Kyurkchiev, N., S. Markov. On the Hausdorff Distance between the Heaviside Step Function and Verhulst Logistic Function. – J. Math. Chem., Vol. **54**, 2016, No 1, pp. 109-119.
24. Kyurkchiev, N., S. Markov. Sigmoid Functions: Some Approximation and Modelling Aspects. Saarbrücken, LAP LAMBERT Academic Publishing, 2015. ISBN 978-3-659-76045-7.
25. Iliev, A., N. Kyurkchiev, A. Rahnev, T. Terzieva. Some Models in the Theory of Computer Viruses Propagation. LAP LAMBERT Academic Publishing, 2019. ISBN: 978-620-0-00826-8.
26. Markov, S. Reaction Networks Reveal New Links between Gompertz and Verhulst Growth Functions. – Biomath, Vol. **8**, 2019, No 1.
27. Kyurkchiev, N. On a Sigmoidal Growth Function Generated by Reaction Networks. Some Extensions and Applications. – Communications in Applied Analysis, Vol. **23**, 2019, No 3, pp. 383-400.
28. Kyurkchiev, N., A. Iliev, S. Markov. Some Techniques for Recurrence Generating of Activation Functions: Some Modeling and Approximation Aspects. LAP LAMBERT Academic Publishing, 2017. ISBN: 978-3-330-33143-3.
29. Kyurkchiev, N., A. Iliev, A. Rahnev. Some Families of Sigmoid Functions: Applications to Growth Theory. LAP LAMBERT Academic Publishing, 2019. ISBN: 978-613-9-45608-6.
30. Iliev, A., N. Kyurkchiev, S. Markov. A Note on the New Activation Function of Gompertz Type. – Biomath Communications, Vol. **4**, 2017, No 2. 20 p.
31. Anguelov, R., M. Borisov, A. Iliev, N. Kyurkchiev, S. Markov. On the Chemical Meaning of Some Growth Models Possessing Gompertzian-Type Property. – Math. Meth. Appl. Sci., 2017, pp. 1-12.
32. Anguelov, R., N. Kyurkchiev, S. Markov. Some Properties of the Blumberg's Hyper-Log-Logistic Curve. – BIOMATH, Vol. **7**, 2018, No 1. 8 p.
33. Markov, S., A. Iliev, A. Rahnev, N. Kyurkchiev. On the Exponential-Generalized Extended Compertz Cumulative Sigmoid. – International Journal of Pure and Applied Mathematics, Vol. **120**, 2018, No 4, pp. 555-562.
34. Pavlov, N., A. Iliev, A. Rahnev, N. Kyurkchiev. Some Software Reliability Models: Approximation and Modeling Aspects. LAP LAMBERT Academic Publishing, 2018. ISBN: 978-613-9-82805-0.
35. Pavlov, N., A. Iliev, A. Rahnev, N. Kyurkchiev. Nontrivial Models in Debugging Theory. Part 2. LAP LAMBERT Academic Publishing, 2018. ISBN: 978-613-9-87794-2.
36. Iliev, A., N. Kyurkchiev, S. Markov. On the Approximation of the Cut and Step Functions by Logistic and Gompertz Functions. – Biomath, Vol. **4**, 2015, pp. 2-13.
37. Kyurkchiev, N., A. Iliev, A. Rahnev. A New Class of Activation Functions Based on the Correcting Amendments of Gompertz-Makeham Type. – Dynamic Systems and Applications, Vol. **28**, 2019, No 2, pp. 243-257.
38. Kyurkchiev, N. Investigations on a Hyper-Logistic Model. Some Applications. – Dynamic Systems and Applications, Vol. **28**, 2019, No 2, pp. 351-369.
39. Markov, S., A. Iliev, A. Rahnev, N. Kyurkchiev. A Note on the n-Stage Growth Model. Overview. – Biomath Communications, Vol. **5**, 2018, No 2, pp. 79-100.

40. Kyurkchiev, N., A. Iliev, A. Rahnev. A Look at the New Logistic Models with “Polynomial Variable Transfer”. LAP LAMBERT Academic Publishing, 2020. ISBN: 978-620-2-56595-0.
41. Kyurkchiev, N., S. Markov. On a Logistic Differential Model. Some Applications. – Biomath Communications, Vol. **6**, 2019, No 1, pp. 34-50.
42. Kyurkchiev, N. Some New Classes of Growth Functions Generated by Reaction Networks and Based on “Correcting Amendments” of Bateman-Gompertz and Bateman-Gompertz-Makeham-Type, I. – Communications in Applied Analysis, Vol. **24**, 2020, No 1, pp. 13-29.
43. Kyurkchiev, N. On a Class of Growth Curves with Exponentially Variable Transfers Generated by Reaction Networks, II. – International Electronic Journal of Pure and Applied Mathematics, Vol. **14**, 2020, No 1, pp. 21-29.
44. <http://portalms.saude.gov.br/images/pdf/2016/maio/24/Informe-Epidemiol--gico-n---27--SE-20-2016--24mai2016-12h00.pdf>
45. <https://virologydownunder.blogspot.com/2016/06/brazils-microcephaly-and-cns-disorder-m.html>
46. https://www.who.int/csr/disease/swineflu/history_map/InfluenzaAH1N1_maps.html

Received: 05.09.2020; Second Version: 23.10.2020; Accepted: 30.10.2020