

Oceanological and Hydrobiological Studies

International Journal of Oceanography and Hydrobiology

Volume 40, Issue 3

ISSN 1730-413X

eISSN 1897-3191

(28–35)

2011



DOI: 10.2478/s13545-011-0026-3
Original research paper

Received:

January 14, 2011

Accepted:

March 31, 2011

Has zooplankton 24 hour vertical distribution pattern in Lough Derg (Ireland) been changed over the period of ~90 years?

Agnieszka Pociecha^{1,2,*}, Thomas K. McCarthy²

¹ Institute of Nature Conservation, Polish Academy of Sciences, ul. A. Mickiewicza 33, 30-120 Kraków, Poland

² Department of Zoology, National University of Ireland Galway, University Road, Galway, Ireland

Key words: rotifers, crustaceans, vertical distribution, migration, Irish lake

Abstract

The vertical distribution of zooplankton was examined in Lough Derg, Ireland. Zooplankton was collected at one location every 2 m from the surface to the bottom during 24 hours at four-hour intervals. Zooplankton was identified to the species level and its density was calculated for each taxon, depth and hour. We recorded 9 rotifer, 3 copepod, and 4 cladoceran species. The highest total zooplankton density (rotifers, copepods, cladocerans, mysids and zebra mussel larvae) was recorded at 3 a.m. Rotifers preferred mainly a depth from 0 to 8 m, while copepods and cladocerans were observed within the whole water column during a 24 hour observation. It was a different pattern of diurnal migration than that Southern and Gardiner (1932) received, though they didn't study rotifers. Probably food concentration and/or predators, or other environmental factors could influence the diel vertical migration of zooplankton from Lough Derg.

INTRODUCTION

Heterogeneous distributions of zooplankton originate from various processes, some of which can be attributed to swimming of the animals (Folt, Burns 1999), diel vertical and horizontal migrations (Stich, Lampert 1981; Kvam, Kleiven 1995), and active swimming along horizontal and vertical gradients of temperature, food concentration, predation (Kvam, Kleiven 1995; Lampert et al. 2003), and the abundance of other species (Dumont 1972).

In addition, zooplankton species typically exhibit daily, rhythmic changes in their vertical distribution. In general, zooplankton tend to move upwards during the hours of darkness and downwards as daylight approaches (Southern, Gardiner 1932; Hutchinson 1967).

Lough Derg was chosen to study a 24 hour cycle of zooplankton migration. In the early 1920s, Southern and Gardiner (1932) examined diurnal migrations of crustaceans in Lough Derg and found that species were not homogenous in their pattern of diurnal migration. Since Southern and Gardiner's (1932) study, the lake environment has changed. In subsequent decades, an increase in the abundance of planktonic algae (especially blue-green algae) has been observed, an invasive species – the zebra mussel *Dreissena polymorpha* appeared, and there has been a decrease in the abundance of an important economic fish species – the eel (*Anguilla anguilla*). Further increases in the levels of algal development have been observed in recent years, giving rise to concern about angling and other tourism along the lake shore (McCarthy et al. 1998).

The aim of this study was to re-examine the abundance and species composition of zooplankton across different water layers over a 24 hour period and compare present patterns in zooplankton vertical

* Corresponding author: pociecha@iop.krakow.pl

distribution with those described by Southern and Gardiner (1932).

MATERIALS AND METHODS

Lough Derg (52°49'–53°06' N, 8°09'–8°31' W), the lower of the two larger lakes on the River Shannon is a well mixed, non-freezing, meso- to eutrophic, monomictic lake. Thermocline formation in summer is usually short-lived. Table 1 presents some characteristic features of Lough Derg. Much of the lake lies on carboniferous limestone with the exception of the narrow southern section, which is underlain by Silurian strata. The lower reaches of Lough Derg are largely enclosed by hills, the Slieve Aughty Mountains to the west and the Arra Mountains to the east, while to the north the upper reaches of Lough Derg are surrounded by a relatively flat agricultural landscape (Fig. 1).

Table1

The characteristic features of Lough Derg.

Features	Values
Mean depth (m)	7.6
Max depth (m)	34
Annual water level fluctuation (m)	0.3
Residence time (year)	0.2
Catchment area (km ²)	10.280
Shoreline (km)	179

Bowman (2000) reported some physical and chemical features of the lake during 1998 and 1999: the temperature range from 4.8°C (January) to 17.5°C (August); Secchi disc - 1.0 – 3.5 m; pH - 7.94 – 8.34; chlorophyll - 1 – 12 ($\mu\text{g dm}^{-3}$); N-NO₂ + N-NO₃ - 500 – 1200 ($\mu\text{g dm}^{-3}$); P-PO₄ - 10 – 50 ($\mu\text{g dm}^{-3}$).

Phytoplankton is characterized by diatoms and cyanobacteria. In winter - *Aphanizomenon flos-aquae*, *Oscillatoria* spp., *Stephanodiscus astraea*; in spring - *S. astraea*, *Tabellaria* spp., *Melosira italica*, *Asterionella formosa*; in summer - *Oscillatoria* spp., *Tabellaria* spp., *Melosira granulata*; in autumn - *Oscillatoria* spp., *Tabellaria* spp., *M. granulata* (Bowman 1985, 2000; Bowman et al. 1993). The fish are represented by the following species: *Salmo salar*, *S. trutta*, *Percia fluviatilis*, *Tinca tinca*, *Esox lucius*, *Abramis brama*, *Scardinus erythrophthalmus*, *Anguilla anguilla* (economically important), *Coregonus pollans*, *Rutilus rutilus*. Also an invasive bivalve species, *Dreissena polymorpha*, has been




Fig. 1. Location of Lough Derg (Ireland).

observed in the lake since 1997 (McCarthy et al. 1998).

Samples were taken from the central site at the deepest part of Lough Derg, on 26–27 of August 2003, at four-hour intervals (7 p.m., 11 p.m., 3 a.m., 7 a.m., 11 a.m., 3 p.m.), every two meters (from 0 m to 18 m) (Fig. 1). Samples for zooplankton taxonomic identification and quantitative analyses were collected using a 5-l Ruttner sampler. Ten-liter zooplankton samples (two replicate 5 l samples) were concentrated with a 50 μm plankton net in the field and immediately treated with 4% formalin. For identification and counting of zooplankton species, 5 replicate sub-samples were analyzed microscopically ($\times 100$ or $\times 200$) in the Kolkwitz chamber of 0.4 mm height and 22 mm diameter. Taxonomic analyses of zooplankton (rotifer, copepod and cladoceran taxa) were conducted using the following keys: Dussart (1967, 1969); Flößner (1972, 2000); Ejsmont-Karabin et al. (2004); Rybak and Błędzki (2005); Voight and Koste (1978a,b). Quantitative samples were prepared by filtering 10 l⁻¹ of water and reducing the sample volume to 0.05 l⁻¹. Zooplankters were counted in 0.5 ml⁻¹ Kolkwitz chambers. Mean density per l⁻¹ was estimated on the basis of five chamber counts.

When collecting the water samples for zooplankton community identification, on-site measurements of water temperature ($^{\circ}\text{C}$), pH and conductivity ($\mu\text{S cm}^{-3}$) were also measured using a multi-parameter YSI meter.

RESULTS AND DISCUSSION

During the 24 hour cycle, values of physico-chemical parameters of lake water were similar from surface to bottom, and a thermocline was not present. The temperature ranged between 19.01°C (at 11 a.m.) to 19.25°C (at 7 p.m.), pH was stable at 8.78, and conductivity was $370 \mu\text{S cm}^{-3}$. Similar temperatures, higher pH values and conductivity were recorded in the study of Southern and Gardiner (1926a,b). The lake's high alkalinity is derived from the underlying carboniferous limestone bedrock and run-off from the surrounding Midlands limestone plain.

The ecological importance of migratory behavior of plankton has been the subject of much research and discussion (e.g. Southern, Gardiner 1926a,b; Hutchinson 1967; Dumont et al. 1985). Hutchinson (1967) showed that the maximum zooplankton concentration usually occurs near the surface at night (nocturnal ascent) and at various depths during daytime (diurnal descent). But this is a general point of view and it is likely that many environmental factors could change this "pattern". Southern and Gardiner (1926a,b) in their studies on zooplankton in Lough Derg showed that species were not identical in their reactions to whatever stimuli that provoke the responses of species named "diurnal migration".

During our 24 hour study in Lough Derg, 17 species of zooplankton were found: 9 rotifers, 3 copepods, and 4 cladocerans. Some of them, such as *Keratella cochlearis* and *Polyarthra vulgaris*, are common in the European lakes. One species *Mysis relicta* is typical for water bodies in northern Europe (Appendix 1).

The highest values of total zooplankton density (rotifers, copepods, cladocerans, mysids and zebra mussel larvae) were observed in the water column between 0 and 8 m at 3 a.m. ($780 - 1180 \text{ ind. l}^{-1}$). Zooplankton densities below 100 ind. l^{-1} were observed at 11 a.m. (except at 12 m) and 3 p.m. (except 0 – 4 m). During the early (7 a.m.) and late morning (11 a.m.) the highest densities of zooplankton were observed from 10 m to the bottom but, by the early afternoon (3 p.m.) at 0 to 4 m (Fig. 2). Water-column zooplankton densities were the highest from the surface to 8 m in the late afternoon (7 p.m.), from the surface to 4 m in the

Appendix 1

List of zooplankton species in Lough Derg.

Rotatoria
<i>Ascomorpha saltans</i> Bartsch 1870
<i>Conochilus unicornis</i> Rousselet, 1892
<i>Kellicotia longispina</i> (Kellicott, 1879)
<i>Keratella cochlearis</i> (Gosse, 1879)
<i>Keratella cochlearis</i> f. <i>tecta</i> (Lauterborn, 1900)
<i>Keratella quadrata</i> (Müller, 1786)
<i>Polyarthra vulgaris</i> Carlin, 1943
<i>Synchaeta oblonga</i> Ehrenberg, 1831
<i>Trichocerca similis</i> (Wierzejski, 1893)
Copepoda
<i>Cyclops strenuus</i> (Fischer, 1851)
<i>Eucyclops macrurus</i> (Sars, 1863)
<i>Eudiaptomus gracilis</i> (Sars, 1863)
Cladocera
<i>Bythotrephes longimanus</i> Leydig, 1860
<i>Daphnia galeata</i> Sars, 1864
<i>Daphnia hyalina</i> Leydig, 1860
<i>Leptodora kindtii</i> (Focke, 1844)
Mysidacea
<i>Mysis relicta</i> Loven, 1869
+ larvae of <i>Dreissena polymorpha</i> (Pallas, 1771)

evening (11 p.m.), and from the surface to 8 m at 3 a.m. In general, these observations of total zooplankton density support Hutchinson's hypothesis (1967) that the maximum zooplankton concentration occurs near the surface at night and at various depths during the day.

Rotifers as a group preferred a depth from 0 to 8 m, except at 11 p.m. when they were also observed at the surface (150 ind. l^{-1}) and at the deeper part of the water layer at 18 m (80 ind. l^{-1}). The highest value of rotifer density (167 ind. l^{-1}) was recorded at 3 a.m. at 6 m. A high percentage of rotifers, above 60% of the total zooplankton density, was recorded at 11 p.m. at 8 m, 12 m, 18 m and at 3 a.m. at 6 m and 18 m (Fig. 3). The most abundant rotifer species were *Keratella cochlearis* and *Polyarthra vulgaris*. Both were observed in the whole water column at night, but during the day they were found in different water layers with low abundance (Table 2). George and Fernando (1970) found that these species can exhibit nocturnal upward migration in the summer season and migrate in the reverse direction in winter. Furthermore, Karabin and Ejsmont-Karabin (2005) suggest that the main factor that triggers off the vertical migrations of Rotifera is food resource. However, the presence of predators namely their kairomones may induce vertical migrations (Loose, Dawidowicz 1994)




Fig. 2. Total density (ind. l⁻¹) of zooplankton in a water column of the Lough Derg during the 24 hour study.




Fig. 3. Density of rotifers, copepods and cladocerans in a water column of the Lough Derg during the 24 hour study.

Table 2

Density (ind. l⁻¹) of most abundant species of rotifer and crustacean community during 24 hour cycle in Lough Derg.

Hours	Meters	Rotifera		Copepoda			Cladocera		Others
		<i>Keratella cochlearis</i>	<i>Polyarthra vulgaris</i>	nauplii	copepodite	<i>Eudiaptomus gracilis</i>	<i>Daphnia galeata</i>	<i>Daphnia hyalina</i>	
7 p.m.	0	43		43		51			72
	2	6		45	22	17	17		122
	4	5	19	42		5			154
	6	42	12	79	12	12	24		140
	8	56	6	45		11			106
	10	7	3	3			7		47
	12	7		10		11	3	3	49
	14	10	5	19				10	63
	16		6	5		11		27	43
	18			30	20	20		30	50
11 p.m.	0	101	20	40		74		61	202
	2	17		68	6	52	17	17	232
	4	14	3	24		30		17	48
	6	14		4		8		7	67
	8	31	4			4	4	13	79
	10	13	5	16	5	8		21	91
	12	18	10	3		5		8	34
	14		4	4		18		4	32
	16	9		24				19	57
	18	42	14	56					84
3 a.m.	0	83	33	182		42			670
	2	58	17	58		33		25	291
	4	45	60	165	15	30		45	750
	6	28	125		14	28			599
	8	62	49	99	12	12	25		456
	10	35	26	35	17			26	243
	12	27		27	9	18		9	257
	14	37		18	9	18		9	283
	16	16	24	16	8	16	8	48	129
	18	34						17	206
7 a.m.	0	32	9	27		10		9	73
	2			4		13	17	4	30
	4	13	13	13	7	20		20	91
	6			33		7		7	53
	8		5	9				19	65
	10			33	7	20		60	93
	12		6	18	6	12		6	48
	14	5	5		5	10		5	46
	16		6	66	12	24		12	54
	18			20	7	16		7	30
11 a.m.	0	5		29		10		5	44
	2			19		10		10	29
	4		10	10	7	6		7	27
	6					7		7	37
	8	2		14	2	4		6	14
	10	9	9			14		5	23
	12	11	6	50	28	28		6	106
	14		4	15	4	18			
	16		7	27		7		27	27
	18		4	4		4			9
3 p.m.	0	16	5	26		16		10	88
	2		7	17	10	17		7	63
	4	13	4	4	4	17		4	58
	6		2	2	3	4		6	23
	8			10		13		14	55
	10	4	3	14		3			14

Copepods reached their highest density (200 ind. l⁻¹) at 3 a.m. twice, at 0 m and 4 m. This group preferred the depths from 0 to 8 m at 7 p.m., 11 p.m., 3 a.m. and 3 p.m., but deeper water layers from 12 m to the bottom at 7 a.m. and 11 a.m. A high percentage of copepods (>80% of the total zooplankton density) was observed several times in the morning at 7 a.m. and 11 a.m. (Fig. 3). Immature stages of all copepods (nauplii and copepodites) and *Eudiaptomus gracilis* were most abundant. High numbers of all copepods nauplii and *E. gracilis* were recorded throughout the whole water column while copepodites of all copepods were found at specific water layers (Table 2). This differs from the patterns of *E. gracilis* migration observed by Southern and Gardiner (1932). They observed *E. gracilis* adult and immature stages with the maximum concentration in the surface layers (0–5 m) during the night and at 10–20 m during the day. In another Irish lake (Gouganebarra), Gardiner (1957) observed the same pattern of *Diaptomus laticeps* diurnal migration with maximum concentrations at 0–5 m during night time (3 a.m.) and at 20–25 m during daytime (2 p.m.).

Cladoceran density was never above 100 ind. l⁻¹. The highest densities of cladocerans were observed during the night in different water layers: 11 p.m. at 0 m (68 ind. l⁻¹), 3 a.m. at 4 m (75 ind. l⁻¹) and 16 m (64 ind. l⁻¹); and during the early morning at 7 a.m. at 10 m (60 ind. l⁻¹). In general cladocerans preferred surface water layers during the night but deeper water layers during the day. A high percentage of cladocerans, 40–55% of the total density of zooplankton, was recorded at 2 m (7 a.m.), 6 m (11 a.m. and 3 p.m.), between 8–10 m (7 a.m.) and once in a deeper layer at 16 m (11 p.m.) (Fig. 3).

Daphnia hyalina was the most abundant species and was observed within the whole water column except at 7 p.m. when it was found only below 12 m (Table 2). Southern and Gardiner (1932) reported a contrasting pattern of migration for *D. longispina*, a large *Daphnia* species similar to *D. hyalina*. During the morning (6 a.m.) they found high concentrations of *D. longispina* in deeper water layers, but by midday *Daphnia* concentrations were the highest in the upper 5 m of the water column and were almost absent below the 10 m level. In addition, Gardiner (1957) found different results for *D. longispina* abundance in Gouganebarra Lake: this species remained at 0–15 m throughout a 24 hour period. Ziarek et al. (2011) observed that individual *Daphnia* behavior shows extreme flexibility as a response to external and

internal factors (food, light, individual size of daphnids, the presence of kairomones, the occurrence of toxic elements), but the most important factor influencing the biology of *Daphnia* is the temperature.

In the present study, *D. polymorpha* larvae were observed in large numbers throughout the whole water column over the 24 hour period examined. The maximum concentration (750 ind. l⁻¹) was observed at 3 a.m. at 4 m depth. High concentrations of *D. polymorpha* larvae (40 to 50% of the total zooplankton community) were commonly observed in Lough Derg during summer months (Pociecha, unpublished data).

The results of our field survey reveal the highly dynamic character and variability of zooplankton behavior. In general, zooplankton vertical migration in Lough Derg was supportive of Hutchinson's (1967) hypothesis. Hutchinson (1967) distinguished three types of vertical migration patterns: nocturnal migration, twilight migration and reverse migration.

We can't distinguished one type of migration for one group in the examined zooplankton groups of Lough Derg. For copepods we can describe nocturnal migration (between 7 p.m. to 3 a.m.) and reverse migration (at 3 p.m.). Cladocerans and rotifers have not any of the typical discussed patterns (Fig. 3). The occurrence of these patterns could have been caused by temperature and light gradients, predator avoidance or presence of invasive species.

The vertical distribution of limnetic zooplankton species in Lough Derg may be related to the absence of a thermocline as a borderline for zooplankton species (Lampert 2005). Temperature is often observed to be the most important factor governing the zooplankton vertical distribution (Dawidowicz, Loose 1992; Cooke et al. 2008). Zooplankton distribution in Lough Derg is probably not connected with homogenous physico-chemical variables but with other environmental factors. Light, temperature and biotic interactions, such as predation, interspecific competition, food availability and feeding habits can influence and modify migration patterns (Gliwicz, Pijanowska 1988; Lampert 1989; Mavuti 1992; Loose, Dawidowicz 1994; Sakwińska, Dawidowicz 2005; Haupt et al. 2009; Cohen, Forward 2009; Wojtal-Frankiewicz et al. 2010). Dumont et al. (1985) stressed that in the absence of strong forcing factors, such as visual predation and light damage, it would be advantageous for zooplankton not to migrate

downwards during the day, allowing zooplankters to feed continuously on the phytoplankton.

Summing up, we did not observe any evidence for a pattern of zooplankton vertical distribution in Lough Derg. Southern and Gardiner (1932) in their studies on the diel vertical migration of crustaceans in Lough Derg didn't obtain a homogenous pattern of DVM, about ~90 years later we stated that crustaceans were observed in the whole water column during day and night. Rotifers preferred mainly a depth from 0 to 8 m but we could not compare this result with the studies of Southern and Gardiner because they investigated only crustaceans. Probably light, food and/or predators, or other environmental factors (especially the concentration of a chemical trigger – the substance kairomone) influence the zooplankton density, but this needs to be confirmed in the future studies. In the future studies we would also like to focus on the pattern of vertical distribution and diel migration of zooplankton in relation to:

- 1) seasons (spring, summer, autumn and winter),
- 2) gradients of light and temperature,
- 3) biotic factors: food availability and predation,
- 4) influence of the invasive species zebra mussel *Dreissena polymorpha*.

Liu et al. 2006 wrote that DVM is only a tip of an iceberg from the perspective of the whole picture of zooplankton behaviors, though it appears to be the most marvelous spectacle during the lifetime of the performers. DVM is notable for its diel rhythm, magnitude of migration and ontogenetic variances (McLaren 1963; Uye et al. 1990; Zaret, Suffern 1976), but all these remarkable features do not mean to decouple the mechanisms underlying DVM from the behavioral strategies directing the “everyday concerns” of an individual animal. In other words, one mechanism, if it is reasonable for interpreting the general behaviors, will be recommendable for the understanding of DVM, and vice versa. Zooplankters seemingly do not need to develop a different logic to deal with the same problem: how to survive as long as possible.

ACKNOWLEDGEMENTS

We would like to thank Prof. Henri Dumont for comments and M.J. Polito and M. Szul for language correction and editorial comments. This study was co-funded by The Ministry of Science and Higher Education in the Ministry's Programme “Support of international researcher mobility” No. 3/MOB/2007/0

and supported by the National University of Ireland, Galway.

REFERENCES

- Bowman J.J., 1985, *The trophic status of Irish lakes: Lough Derg*. An Foras Forbatha, Dublin
- Bowman J.J., 2000, *River Shannon. Lake water quality monitoring 1998 and 1999*. Environmental Protection Agency, Wexford, Ireland, 58-64
- Bowman J.J., McGarigle M. L., Clabby K.I., 1993, *Lough Derg: An investigation of eutrophication and its causes*. A report to the Lough Derg Working Group. Environmental Research Unit, Dublin
- Cohen J.H., Forward R.B., 2009, *Zooplankton diel vertical migration – A review of proximate control*. Oceanography and Marine Biology: An Annual Review, 47: 77-110
- Cooke S.L., Williamson C.E., Leech D.M., Boeing W.J., Torres L., 2008, *Effects of temperature and ultraviolet radiation on diel vertical migration of freshwater crustacean zooplankton*. Canadian Journal of Fisheries and Aquatic Science, 65: 1144-1152
- Dawidowicz, P., Loose, C.J., 1992, *Metabolic costs during predator-induced diel vertical migration of Daphnia*. Limnology and Oceanography, 37: 1589-1595
- Dumont H.J., 1972, *A competition-based approach of the reverse vertical migration in zooplankton and its implications, chiefly based on a study of the interactions of the Rotifer Asplanchna priodonta (Gosse) with several Crustacea Entomostraca*. Internationale Revue der Gesamten Hydrobiologie und Hydrographie, 57: 1-38
- Dumont H.J., Guisez Y., Carels I., Verheye H.M., 1985, *Experimental isolation of positively and negatively phototactic phenotypes from a natural population of Daphnia magna Strauss: A contribution to the genetic of vertical migration*. Hydrobiologia, 126: 121-127
- Dussart B., 1967, *Les Copépodes des eaux continentales d'Europe occidentale: Calanoides et harpacticopides*. Paris: Boubée and Cie, pp. 500
- Dussart B., 1969, *Les Copépodes des eaux continentales d'Europe occidentale: Cyclopoides et biologie*. Paris: Boubée and Cie, pp. 267
- Ejsmont-Karabin J., Radwan S. and Bielańska-Grainier I., 2004, *Polish Freshwater Fauna. Rotifers (Rotifera)*. Łódź, University of Łódź, pp. 447
- Flößner D., 1972, *Krebstiere, Crustacea. Kiemen-und Blattfüßer, Branchiopoda. Fischläuse, Branchiura*. Jena, Fischer Verlag, pp. 501
- Flößner D., 2000, *Die Haplopoda und Cladocera Mitteleuropas*. Leiden, Backhuys Publisher, pp. 425
- Folt C.L., Burns C.W., 1999, *Biological drivers of zooplankton patchiness*. Trends in Ecology & Evolution, 14: 300-305
- Gardiner A.C., 1957, *Preliminary observations on the diurnal migration of the Crustacea in the plankton of Gonganebarra Lake*. Proceedings of the Royal Irish Academy, 58B: 305-319
- George M.F., Fernando C.H., 1970, *Diurnal migration in three species of rotifers in Sunfish Lake, Ontario*. Limnology and Oceanography, 15: 218-223
- Gliwicz M.Z., Pijanowska J., 1988, *Effect of predation and resource depth distribution on vertical migration of zooplankton*. Bulletin of Marine Science, 43: 695-709
- Haupt F., Stockenreiter M., Baumgartner M., Boersma M., Stibor H., 2009, *Daphnia diel vertical migration: implications beyond zooplankton*. Journal of Plankton Research, 31: 515-524
- Hutchinson G. E., 1967, *A treatise on Limnology, vol II: Introduction to Lake Biology and the Limnoplankton*. Wiley, New York, pp. 1115

- Karabin A., Ejsmont-Karabin J., 2005, *An evidence for vertical migrations of small rotifers – a case of rotifer community in a dystrophic lake*. Hydrobiologia, 546: 381–386
- Kvam O.V., Kleiven O.T., 1995, *Diel horizontal migration and swarm formation in Daphnia in response to Chaoborus*. Hydrobiologia, 307: 177–184
- Lampert W., 1989, *The adaptive significance of diel vertical migration of zooplankton*. Functional Ecology, 3: 21–27
- Lampert W., 2005, *Vertical distribution of zooplankton: density dependence and evidence for an ideal free distribution with costs*. BMC Biology, 3:10, 10, doi:10.1186/1741-7007-3-10
- Lampert W., McCauley E., Manly B.F.J., 2003, *Trade-offs in the vertical distribution of zooplankton: ideal free distribution with costs?* Proceedings of the Royal Society of London, Series B, 270: 765–773
- Loose C.J., Dawidowicz P., 1994, *Trade-offs in diel vertical migration by zooplankton: the costs of predator avoidance*. Ecology, 75: 2255–2263
- Liu S-H., Sun S., Han B-P., 2006, *Viewing DVM via general behaviors of zooplankton: A way bridging the success of individual and population*. Journal of Theoretical Biology 238: 435–448
- McCarthy T.K., Fitzgerald J., O'Connor W., 1998, *The occurrence of the zebra mussel Dreissena polymorpha (Pallas. 1771), an introduced biofouling freshwater bivalve in Ireland*. Irish Naturalists' Journal, 25: 413–415
- McLaren I.A., 1963, *Effects of temperature on growth of zooplankton and the adaptive value of vertical migration*. Journal of the Fisheries Research Board of Canada, 26: 199–220
- Mavuti K.M., 1992, *Diel vertical distribution of zooplankton in Lake Naivasha, Kenya*. Hydrobiologia, 232: 31–41
- Rybák J.L., Błędzki L.A., 2005, *Copepods. Copepoda: Cyclopoida*. Warszawa: IOS, BMŚ, pp. 127
- Sakwińska O., Dawidowicz P., 2005, *Life history strategy and depth selection behavior as alternative antipredator defenses among natural Daphnia hyalina populations*. Limnology and Oceanography, 50: 1284–1289
- Southern R., Gardiner A.C., 1926a, *Reports from the Limnological Laboratory. I. The Seasonal Distribution of the Crustacea of the Plankton in Lough Derg and the River Shannon*. Fisheries, Ireland, Science Investment pp. 1–170. Plates I–XV. Text figs. 1–4
- Southern R., Gardiner A.C., 1926b, *A preliminary account of the diurnal migration of the Crustacea of the plankton of Lough Derg*. Internationale Revue der Gesamten Hydrobiologie und Hydrographie, 5/6: 323–326
- Southern R., Gardiner A.C., 1932, *The diurnal migration of the Crustacea of the plankton in Lough Derg*. Proceedings of the Royal Irish Academy, 40B: 121–159
- Stich H.B., Lampert W., 1981, *Predator evasion as an explanation of diurnal vertical migration by zooplankton*. Nature, 293: 396–398
- Voigt M., Koste W., 1978a, *Rotatoria. Die Rädertiere Mitteleuropas. Überordnung Monogononta. I. Textband*. Berlin-Stuttgart, Gebrüder Borntraeger, pp. 673
- Voigt M., Koste W., 1978b, *Rotatoria. Die Rädertiere Mitteleuropas. Überordnung Monogononta. II. Tafelband*. Berlin-Stuttgart, Gebrüder Borntraeger, pp. 468
- Uye S., Huang C., Onbe T., 1990, *Ontogenetic diel vertical migration of the planktonic copepod Calanus sinicus in the Inland Sea of Japan*. Marine Biology, 104: 389–396
- Wojtal-Frankiewicz A., Frankiewicz P., Jurczak T., Grennan J., McCarthy T.K., 2010, *Comparison of fish and phantom midge influence on cladocerans diel vertical migration in a dual basin lake*. Aquatic Ecology, 44: 243–254
- Zaret T.M., Suffern, J.S., 1976, *Vertical migration in zooplankton as a predator avoidance mechanism*. Limnology and Oceanography, 21: 804–813
- Ziarek J.J., Nihongi A., Nagai T., Uttieri M., Strickler J.R., 2011, *Seasonal adaptations of Daphnia pulicaria swimming behaviour: the effect of water temperature*. Hydrobiologia, 661: 317–327