THE DISTRIBUTIONAL ASYMPTOTICS MOD 1 OF $(\log_b n)$

CHUANG XU

University of Copenhagen, Copenhagen, DENMARK

ABSTRACT. This paper studies the distributional asymptotics of the slowly changing sequence of logarithms $(\log_b n)$ with $b \in \mathbb{N} \setminus \{1\}$. It is known that $(\log_b n)$ is not uniformly distributed modulo one, and its omega limit set is composed of a family of translated exponential distributions with constant $\log b$. An improved upper estimate $(\sqrt{\log N}/N)$ is obtained for the rate of convergence with respect to (w.r.t.) the *Kantorovich metric* on the circle, compared to the general results on rates of convergence for a class of slowly changing sequences in the author's companion in-progress work. Moreover, a sharp rate of convergence $(\log N/N)$ w.r.t. the *Kantorovich metric* on the interval $[0, 1]$, is derived. As a byproduct, the rate of convergence w.r.t. the *discrepancy metric* (or the *Kolmogorov metric*) turns out to be $(\log N/N)$ as well, which verifies that an upper bound for this rate derived in [OHKUBO, Y.—STRAUCH, O.: *Distribution of leading digits of numbers*, Unif. Distrib. Theory, **11** (2016), no.1, 23–45.] is sharp.

Communicated by Reinhard Winkler

1. Introduction

Given a sequence of real numbers (x_n) , associate with it a sequence $(\nu_N(x_n))_{N \geq 1}$ of finitely supported probability measures

$$\nu_N(x_n) := \frac{1}{N} \sum_{n=1}^N \delta_{\langle x_n \rangle},$$

© 2019 BOKU-University of Natural Resources and Life Sciences and Mathematical Institute, Slovak Academy of Sciences.

2010 Mathematics Subject Classification: 11K06, 11K31, 60B10, 60E20.

Keywords: Uniformly distributed modulo one sequence, slowly changing sequence, rate of convergence, Kantorovich metric, discrepancy, probability measures.

This research is supported in part by a Pacific Institute for the Mathematical Sciences (PIMS) Graduate Scholarship and a Josephine Mitchell Graduate Scholarship at the University of Alberta.

Licensed under the Creative Commons Attribution-NC-ND 4.0 International Public License.

where $\delta_{\langle x_n \rangle}$ stands for the Dirac measure concentrated at $\langle x_n \rangle$, the natural projection of x_n onto the circle $\mathbb{T} = \mathbb{R}/\mathbb{Z}$. Here and throughout, we write ν_N for $\nu_N(x_n)$ when (x_n) is clear from the context. Note that (ν_N) is a sequence in the space $\mathcal{P}(\mathbb{T})$ of all Borel probability measures on $\mathbb{T}$. As a set, $\mathcal{P}(\mathbb{T})$ can be identified with the subspace $\{\mu \in \mathcal{P}(\mathbb{I}) : \mu(\{1\}) = 0\}$ of $\mathcal{P}(\mathbb{I})$, where $\mathbb{I}$ denotes the compact unit interval $[0, 1]$. Lowercase Greek letters μ, ν are used henceforth to denote elements of both $\mathcal{P}(\mathbb{T})$ and $\mathcal{P}(\mathbb{I})$, but it will always be clear from the context which space of measures is meant. Recall that a sequence (x_n) in $\mathbb{R}$ is *uniformly distributed modulo one* (u.d. mod 1) [6, Ch.1] if ν_N converges weakly in $\mathcal{P}(\mathbb{T})$ to the uniform distribution $\lambda_{\mathbb{T}}$ on $\mathbb{T}$. Denote $\lambda_{\mathbb{I}}$ the uniform distribution on $\mathbb{I}$, and let d_K denote the *discrepancy* (or *Kolmogorov*) metric on $\mathcal{P}(\mathbb{I})$, i.e.,

$$d_K(\mu, \nu) = \sup_{x \in \mathbb{I}} |\mu([0, x]) - \nu([0, x])|, \quad \forall \mu, \nu \in \mathcal{P}(\mathbb{I}).$$

Recall from [6, Cor.2.1.1] that (x_n) is u.d. mod 1 if and only if $\lim_{N \rightarrow \infty} d_K(\nu_N \circ \iota^{-1}, \lambda_{\mathbb{I}}) = 0$, where $\iota : \mathbb{T} \rightarrow \mathbb{I}$ is the natural inclusion; see Section 2 for details. It is well known [6, Cor.2.1.2&Thm.2.2.2] that $d_K(\nu_N \circ \iota^{-1}, \lambda_{\mathbb{I}}) \geq \frac{1}{2N}$ for every positive integer N ; in fact, given any (x_n) there exists a constant $c > 0$ such that $d_K(\nu_N \circ \iota^{-1}, \lambda_{\mathbb{I}}) > c \log N/N$ for infinitely many N .

There is a vast literature on the estimation of discrepancy, especially for u.d. mod 1 sequences. For instance, for the sequence (an) , where $a \in \mathbb{R}$ is irrational with bounded partial quotients, [6, Thm.2.3.4] asserts that

$$d_K(\nu_N \circ \iota^{-1}, \lambda_{\mathbb{I}}) = \mathcal{O}(\log N/N), \quad (1)$$

and (1) also holds for Van der Corput sequence [6, Thm.2.3.5]. However, much less research seems to have been undertaken on sequences that are not u.d. mod 1, for example, on *slowly changing sequences* [5].

Given a sequence (x_n) in $\mathbb{R}$, an improved notion with regard to the distributional asymptotics of x_n is the *Omega limit set* $\Omega[x_n]$, defined as

$$\Omega[x_n] = \{\mu \in \mathcal{P}(\mathbb{T}) : \nu_{N_k}(x_n) \xrightarrow{k \rightarrow \infty} \mu \text{ weakly for some sequence } (N_k) \text{ in } \mathbb{N}\}.$$

It is not hard to see that $\Omega[x_n]$ is non-empty, closed and connected [11]. For sequences (x_n) that are *slowly changing* in the sense that

$$\lim_{n \rightarrow \infty} n(x_{n+1} - x_n) = \xi \in \mathbb{R},$$

it was shown in [5] that (x_n) is not u.d. mod 1; moreover, the elements of $\Omega[x_n]$, have been described in terms of asymptotic distribution functions. Similar results for slowly changing sequences in the literature include logarithms of natural numbers or prime numbers, iterated logarithms, and monotone functions of prime numbers [3, 5–12]. As far as the author knows, however, there were virtually

no results, in the case of slowly changing sequences, on the rate(s) of convergence for subsequences of (ν_N) to $\Omega[x_n]$, not even for very basic sequences such as $(\log_b n)$ with $b \in \mathbb{N} \setminus \{1\}$, prior to [13]. Only recently did the author learn that [9] establishes an upper bound $(\log N/N)$ for the latter, as well as their asymptotic distribution functions. Even there, however, the sharpness of the bound $(\log N/N)$ remains obscure. This article aims at resolving this obscurity. Specifically, for sequences $(\log_b n)$, every limit point in $\Omega[x_n]$ is clearly identified, and $(\log N/N)$ is shown to be the sharp rate of convergence w.r. t. d_K .

While the discrepancy metric (on $\mathcal{P}(\mathbb{T})$, as induced by d_K) has been used in uniform distribution theory for decades, its usage for sequences that are *not* u.d. mod 1 appears debatable. In fact, when analyzing such sequences, it may be more natural to study $\Omega[x_n]$ with a metric metrizing the weak topology of $\mathcal{P}(\mathbb{T})$ such as, for instance, the *Kantorovich* (or *transport*) metric d_T . In a recent work [14], the author obtained several results in this regard, including an upper bound $(\log N/N)$ for d_T -convergence. As is shown in this article, however, this bound is not sharp, and better bounds are provided to replace it for $(x_n) = (\log_b n)$. From the arguments presented, it will also become evident that finding a good *lower* bound remains a formidable challenge, even for sequences as simple as $(\log_b n)$.

2. Preliminaries and notations

Let $\mathbb{R}$, $\mathbb{Z}$, and $\mathbb{N}$ be the set of real numbers, integers, and positive integers, respectively. Recall that $\mathbb{T} = \mathbb{R}/\mathbb{Z}$ can be thought of geometrically as the unit circle $\{e^{2\pi i x} : x \in \mathbb{R}\}$ in the complex plane, with its usual topology. For $-\infty < a < b \leq \infty$, let $[a, b[:= \{y \in \mathbb{R} : a \leq y < b\}$; intervals $[a, b]$, $]a, b]$, $]a, b[$ are defined analogously. Let $\lfloor x \rfloor$, $\lceil x \rceil$, and $\langle\langle x \rangle\rangle = x - \lfloor x \rfloor$ be the floor (i.e., the largest integer $\leq x$), the ceiling (i.e., the smallest integer $\geq x$), and the fractional part of $x \in \mathbb{R}$, respectively. In what follows, it will prove useful to denote by $\pi : \mathbb{R} \rightarrow \mathbb{T}$, with $\pi(x) = \langle x \rangle = x + \mathbb{Z}$, and by $\iota : \mathbb{T} \rightarrow \mathbb{I}$, with $\iota(\langle x \rangle) = \langle\langle x \rangle\rangle$, the natural projection and inclusion, respectively. Arguably the most fundamental maps on $\mathbb{T}$ are rotations: Given any $\theta \in \mathbb{R}$, let R_θ be the (counter-clockwise) rotation of $\mathbb{T}$ by $2\pi\theta$, that is, $R_\theta(\langle x \rangle) = \langle x + \theta \rangle$ for all $\langle x \rangle \in \mathbb{T}$. With this, clearly $R_\theta^k = R_{k\theta} = R_{\langle\langle k\theta \rangle\rangle}$ for all $\theta \in \mathbb{R}$ and $k \in \mathbb{Z}$.

Let (X, ρ_X) be a compact metric space, and $\mathcal{P}(X)$ the space of all Borel probability measures on X , endowed with the weak topology. Recall that $\mathcal{P}(X)$

is compact and metrizable. The *Kantorovich distance* on X is

$$d_X(\mu, \nu) = \inf_{\gamma} \int_{X \times X} \rho_X(x, y) d\gamma(x, y), \quad \forall \mu, \nu \in \mathcal{P}(X),$$

where the infimum is taken over all Borel probability measures γ on $X \times X$ with marginals μ and ν . Note that d_X metrizes the weak topology of $\mathcal{P}_X$. For $X = \mathbb{I}$ and $X = \mathbb{T}$, let $\rho_{\mathbb{I}} = |x - y|$, and $\rho_{\mathbb{T}}(x, y) = \min\{|\iota(x) - \iota(y)|, 1 - |\iota(x) - \iota(y)|\}$, $\forall x, y \in X$, respectively. Note also that $\mu \mapsto \mu \circ \pi^{-1}$ maps $\mathcal{P}(\mathbb{I})$ continuously onto $\mathcal{P}(\mathbb{T})$; when restricted to $\tilde{\mathcal{P}}(\mathbb{I}) := \{\mu \in \mathcal{P}(\mathbb{I}) : \mu(\{1\}) = 0\}$, a dense G_{δ} -set in $\mathcal{P}(\mathbb{I})$, this even yields a continuous bijection, but not a homeomorphism, as $\tilde{\mathcal{P}}(\mathbb{I})$ is not compact. In the opposite direction, $\mu \mapsto \mu \circ \iota^{-1}$ establishes a measurable bijection from $\mathcal{P}(\mathbb{T})$ onto $\tilde{\mathcal{P}}(\mathbb{I})$. Note also that $\mu \mapsto \mu \circ R_{\theta}^{-1}$ defines a homeomorphism of $\mathcal{P}(\mathbb{T})$.

Recall that $d_{\mathbb{I}}$ can be expressed explicitly as

$$d_{\mathbb{I}}(\mu, \nu) = \int_0^1 |F_{\mu}(x) - F_{\nu}(x)| dx, \quad \forall \mu, \nu \in \mathcal{P}(\mathbb{I}), \quad (2)$$

where F_{ν} is the distribution function of ν . A method of computing $d_{\mathbb{T}}$ has been developed in [1]; only the following simple upper bound will be used here.

PROPOSITION 2.1. [1, Cor.3.8]. *Assume that $\mu, \nu \in \mathcal{P}(\mathbb{T})$. Then*

$$\begin{aligned} d_{\mathbb{T}}(\mu, \nu) &\leq \inf_{y \in \mathbb{I}} \int_0^1 |(F_{\mu \circ \iota^{-1}}(x) - F_{\nu \circ \iota^{-1}}(x)) - (F_{\mu \circ \iota^{-1}}(y) - F_{\nu \circ \iota^{-1}}(y))| dx \\ &\leq d_K(\mu \circ \iota^{-1}, \nu \circ \iota^{-1}). \end{aligned}$$

For every $a > 0$, consider the negative exponential distribution $-\text{Exp}(a)$ on $\mathbb{R}$ with parameter a , that is,

$$F_{-\text{Exp}(a)}(x) = e^{ax}, \quad \forall x \leq 0,$$

and let $E_a = -\text{Exp}(a) \circ \pi^{-1} \in \mathcal{P}(\mathbb{T})$. Thus

$$F_{E_a \circ \iota^{-1}}(x) = \frac{e^{ax} - 1}{e^a - 1}, \quad \forall x \in \mathbb{I}.$$

Rotated versions of E_a , that is, probabilities $E_a \circ R_{\theta}^{-1}$ with $\theta \in \mathbb{R}$, play an important role in this article. For such probabilities, observe that

$$F_{E_a \circ R_{\theta}^{-1} \circ \iota^{-1}} = \begin{cases} \frac{e^{\langle\langle\theta\rangle\rangle(e^{ax}-1)}}{e^a-1} & \text{if } x \in [0, 1 - \langle\langle\theta\rangle\rangle], \\ 1 + \frac{e^{\langle\langle\theta\rangle\rangle(e^{a(x-1)}-1)}}{e^a-1} & \text{if } x \in [1 - \langle\langle\theta\rangle\rangle, 1[. \end{cases}$$

Henceforth, our analysis focuses on the sequences $(x_n) = (\log_b n)$ with $b \in \mathbb{N} \setminus \{1\}$, and the associated discrete measures $\nu_N = \nu_N(\log_b n) \in \mathcal{P}(\mathbb{T})$. A simple calculation yields an explicit formula for the distribution function of $\nu_N \circ \iota^{-1}$.

PROPOSITION 2.2. *Assume that $b \in \mathbb{N} \setminus \{1\}$ and $N \in \mathbb{N}$. Then, with*

$$L = \lfloor \log_b N \rfloor,$$

$$F_{\nu_N \circ \iota^{-1}}(x) = \begin{cases} \frac{L+1+\sum_{j=0}^L (\lfloor ib^{-j} \rfloor - b^{L-j})}{N} & \text{if } x \in \left[\log_b \frac{i}{b^L}, \log_b \frac{i+1}{b^L} \right], \\ & i = b^L, \dots, N-1, \\ 1 + \frac{L+1+\sum_{j=0}^L (\lfloor \lfloor Nb^{-1} \rfloor b^{-j} \rfloor - b^{L-j})}{N} & \text{if } x \in \left[\log_b \frac{N}{b^L}, \log_b \frac{b \lfloor Nb^{-1} \rfloor + b}{b^L} \right], \\ 1 + \frac{L+1+\sum_{j=0}^L (\lfloor ib^{-j} \rfloor - b^{L-j})}{N} & \text{if } x \in \left[\log_b \frac{bi}{b^L}, \log_b \frac{b(i+1)}{b^L} \right], \\ & i = \lfloor Nb^{-1} \rfloor + 1, \dots, b^L - 1. \end{cases} \quad (3)$$

3. Rates of convergence

In this section, we study the rate of convergence for subsequences of

$$(\nu_N)_{N \geq 1} \text{ w.r.t. } d_{\mathbb{T}}, d_{\mathbb{I}}, \text{ and } d_K.$$

Throughout, for ease of exposition, all proofs are given for $b = 10$, but all arguments can easily be adjusted to any other base $b \in \mathbb{N} \setminus \{1\}$.

3.1. Upper bound for the rate of convergence w.r.t. $d_{\mathbb{T}}$

We first present our main result regarding an upper bound for the rate of convergence w.r.t. $d_{\mathbb{T}}$.

THEOREM 3.1. *Assume $b \in \mathbb{N} \setminus \{1\}$. Then*

$$\limsup_{N \rightarrow \infty} \frac{N}{\sqrt{\log N}} d_{\mathbb{T}} \left(\nu_N, E_{\log b} \circ R_{-\log_b N}^{-1} \right) < +\infty. \quad (4)$$

Proof. Recall that $b = 10$ is assumed throughout; for every $N \in \mathbb{N}$, let $n = \lfloor \log_{10} N \rfloor + 1$ for convenience, and thus $10^{n-1} \leq N \leq 10^n - 1$, as well as

$$\eta_N = E_{\log_b N} \circ R_{-\log_b N}^{-1} \in \mathcal{P}(\mathbb{T}).$$

By Proposition 2.1, it suffices to estimate

$$\int_0^1 \left| (F_{\nu_{N \circ \ell^{-1}}}(x) - F_{\eta_{N \circ \ell^{-1}}}(x)) - (F_{\nu_{N \circ \ell^{-1}}}(y) - F_{\eta_{N \circ \ell^{-1}}}(y)) \right| dx,$$

for an appropriate $0 \leq y < 1$.

Utilizing Proposition 2.2 we first simplify the latter expression as follows: For every $y \in [0, \log_{10} N - n + 1[$, let $i_0 = \lfloor 10^{y+n-1} \rfloor$. Then $10^{n-1} \leq i_0 \leq N-1$ and $y \in [\log_{10} i_0 - n + 1, \log_{10}(i_0 + 1) - n + 1[$. Similarly, for $10^{n-1} \leq i \leq 10^n - 1$ and $x \in \mathbb{I}$,

$$x \in [\log_{10} i - n + 1, \log_{10}(i + 1) - n + 1[\Leftrightarrow i = \lfloor 10^{x+n-1} \rfloor. \quad (5)$$

With this, it follows from Proposition 2.2 that

$$\begin{aligned} & \int_0^1 \left| (F_{\nu_{N \circ \ell^{-1}}}(x) - F_{\eta_{N \circ \ell^{-1}}}(x)) - (F_{\nu_{N \circ \ell^{-1}}}(y) - F_{\eta_{N \circ \ell^{-1}}}(y)) \right| dx \\ &= \sum_{i=10^{n-1}}^{N-1} \int_{\log_{10} i - n + 1}^{\log_{10}(i+1) - n + 1} \left| \frac{\sum_{j=0}^{n-1} (\lfloor i 10^{-j} \rfloor - \lfloor i_0 10^{-j} \rfloor)}{N} - \frac{10^n (10^x - 10^y)}{9N} \right| dx \\ &+ \sum_{i=\lfloor N/10 \rfloor + 1}^{\log_{10}(\lfloor N/10 \rfloor + 1) - n + 2} \int_{\log_{10}(N/10) - n + 2}^{\log_{10}(i+1) - n + 2} \left| \frac{\sum_{j=0}^{n-1} (\lfloor \lfloor N/10 \rfloor 10^{-j} \rfloor - \lfloor i_0 10^{-j} \rfloor)}{N} - \frac{10^n (10^{x-1} - 10^y)}{9N} \right| dx \\ &+ \sum_{i=\lfloor N/10 \rfloor + 1}^{10^{n-1}-1} \int_{\log_{10} i - n + 2}^{\log_{10}(i+1) - n + 2} \left| \frac{\sum_{j=0}^{n-1} (\lfloor i 10^{-j} \rfloor - \lfloor i_0 10^{-j} \rfloor)}{N} - \frac{10^n (10^{x-1} - 10^y)}{9N} \right| dx \\ &= \sum_{i=\lfloor N/10 \rfloor + 1}^{N-1} \int_{\log_{10} i - n + 1}^{\log_{10}(i+1) - n + 1} \left| \frac{\sum_{j=0}^{n-1} (\lfloor i 10^{-j} \rfloor - \lfloor i_0 10^{-j} \rfloor)}{N} - \frac{10^n (10^x - 10^y)}{9N} \right| dx \end{aligned}$$

$$+ \int_{\log_{10}(N/10)-n+1}^{\log_{10}(\lfloor N/10 \rfloor + 1) - n + 1} \left| \frac{\sum_{j=0}^{n-1} (\lfloor \lfloor N/10 \rfloor 10^{-j} \rfloor - \lfloor i_0 10^{-j} \rfloor)}{N} - \frac{10^n(10^x - 10^y)}{9N} \right| dx.$$

Since $|F_{\nu_N \circ \iota^{-1}}(x) - F_{\eta_N \circ \iota^{-1}}(x)| \leq 1$ for all $x \in \mathbb{I}$, it easily follows that

$$\int_{\log_{10}(N/10)-n+1}^{\log_{10}(\lfloor N/10 \rfloor + 1) - n + 1} \left| \frac{\sum_{j=0}^{n-1} (\lfloor \lfloor N/10 \rfloor 10^{-j} \rfloor - \lfloor i_0 10^{-j} \rfloor)}{N} - \frac{10^n(10^x - 10^y)}{9N} \right| dx = \mathcal{O}(N^{-1}).$$

From (5) and $i_0 = \lfloor 10^{y+n-1} \rfloor$, it is readily verified that

$$\begin{aligned} & \frac{\sum_{j=0}^{n-1} (\lfloor i 10^{-j} \rfloor - \lfloor i_0 10^{-j} \rfloor)}{N} - \frac{10^n(10^x - 10^y)}{9N} \\ &= \frac{\sum_{j=0}^{n-1} ((i_0 10^{-j} - \lfloor i_0 10^{-j} \rfloor) - (i 10^{-j} - \lfloor i 10^{-j} \rfloor))}{N} \\ &+ \frac{10((\lfloor 10^{x+n-1} \rfloor - 10^{x+n-1}) - (\lfloor 10^{y+n-1} \rfloor - 10^{y+n-1}))}{9N} \\ &- \frac{10^{1-n}(\lfloor 10^{x+n-1} \rfloor - \lfloor 10^{y+n-1} \rfloor)}{9N}. \end{aligned}$$

From

$$\left| \frac{10((\lfloor 10^{x+n-1} \rfloor - 10^{x+n-1}) - (\lfloor 10^{y+n-1} \rfloor - 10^{y+n-1}))}{9N} - \frac{10^{1-n}(\lfloor 10^{x+n-1} \rfloor - \lfloor 10^{y+n-1} \rfloor)}{9N} \right| = \mathcal{O}(N^{-1}),$$

for $y \in [0, \log_{10} N - n + 1[$, it follows that

$$\begin{aligned} & \int_0^1 |(F_{\nu_N \circ \iota^{-1}}(x) - F_{\eta_N \circ \iota^{-1}}(x)) - (F_{\nu_N \circ \iota^{-1}}(y) - F_{\eta_N \circ \iota^{-1}}(y))| dx = \\ & \frac{1}{N} \sum_{i=\lfloor N/10 \rfloor + 1}^{N-1} \log_{10} \left(1 + \frac{1}{i} \right) \cdot \left| \sum_{j=0}^{n-1} ((i_0 10^{-j} - \lfloor i_0 10^{-j} \rfloor) - (i 10^{-j} - \lfloor i 10^{-j} \rfloor)) \right| + \\ & \mathcal{O}(N^{-1}). \quad (6) \end{aligned}$$

In the following, we further estimate the right hand side of (6). The elementary inequality

$$x - x^2/2 \leq \log(x+1) \leq x, \quad \forall x \geq 0$$

yields

$$\frac{1}{i \log 10} - \frac{1}{2i^2 \log 10} \leq \log_{10} \left(1 + \frac{1}{i} \right) \leq \frac{1}{i \log 10};$$

moreover,

$$\frac{1}{N} \sum_{i=\lfloor N/10 \rfloor + 1}^{N-1} \frac{1}{2i^2 \log 10} \left| \sum_{j=0}^{n-1} ((i_0 10^{-j} - \lfloor i_0 10^{-j} \rfloor) - (i 10^{-j} - \lfloor i 10^{-j} \rfloor)) \right| = \mathcal{O}(N^{-2} \log N).$$

Hence

$$\begin{aligned} & \int_0^1 \left| (F_{\nu_{N \circ \ell^{-1}}}(x) - F_{\eta_{N \circ \ell^{-1}}}(x)) - (F_{\nu_{N \circ \ell^{-1}}}(y) - F_{\eta_{N \circ \ell^{-1}}}(y)) \right| dx = \\ & \frac{1}{N \log 10} \sum_{i=\lfloor N/10 \rfloor + 1}^{N-1} \frac{1}{i} \left| \sum_{j=0}^{n-1} (i_0 10^{-j} - \lfloor i_0 10^{-j} \rfloor) - (i 10^{-j} - \lfloor i 10^{-j} \rfloor) \right| + \\ & \mathcal{O}(N^{-1}). \end{aligned}$$

From

$$\begin{aligned} & \frac{1}{N \log 10} \frac{1}{i} \left| \sum_{j=0}^{n-1} ((i_0 10^{-j} - \lfloor i_0 10^{-j} \rfloor) - (i 10^{-j} - \lfloor i 10^{-j} \rfloor)) \right| \\ & \leq \frac{2n}{N \lfloor N/10 \rfloor \log 10}, \quad \text{for } i = \lfloor N/10 \rfloor \quad \text{and} \quad i = N, \end{aligned}$$

it follows that

$$\begin{aligned} & \int_0^1 \left| (F_{\nu_{N \circ \ell^{-1}}}(x) - F_{\eta_{N \circ \ell^{-1}}}(x)) - (F_{\nu_{N \circ \ell^{-1}}}(y) - F_{\eta_{N \circ \ell^{-1}}}(y)) \right| dx = \\ & \frac{1}{N \log 10} \sum_{i=\lfloor N/10 \rfloor}^N \frac{1}{i} \left| \sum_{j=0}^{n-1} ((i_0 10^{-j} - \lfloor i_0 10^{-j} \rfloor) - (i 10^{-j} - \lfloor i 10^{-j} \rfloor)) \right| + \\ & \mathcal{O}(N^{-1}). \quad (7) \end{aligned}$$

Completely analogous arguments show that (7) holds also for

$$y \in [\log_{10} N - n + 1, 1[\quad \text{with} \quad i_0 = \lfloor 10^{y+n-2} \rfloor.$$

Thus it suffices to determine the constant order of amplitude of

$$\sum_{i=\lfloor N/10 \rfloor}^N \frac{1}{i} \left| \sum_{j=0}^{n-1} ((i_0 10^{-j} - \lfloor i_0 10^{-j} \rfloor) - (i 10^{-j} - \lfloor i 10^{-j} \rfloor)) \right|.$$

To get rid of the absolute value, one can use the Cauchy-Schwarz inequality:

$$\begin{aligned} & \left\{ \sum_{i=\lfloor N/10 \rfloor}^N \frac{1}{i} \left| \sum_{j=0}^{n-1} ((i_0 10^{-j} - \lfloor i_0 10^{-j} \rfloor) - (i 10^{-j} - \lfloor i 10^{-j} \rfloor)) \right| \right\}^2 \\ & \leq \sum_{i=\lfloor N/10 \rfloor}^N \frac{1}{i^2} \sum_{i=\lfloor N/10 \rfloor}^N \left\{ \sum_{j=0}^{n-1} ((i_0 10^{-j} - \lfloor i_0 10^{-j} \rfloor) - (i 10^{-j} - \lfloor i 10^{-j} \rfloor)) \right\}^2. \end{aligned} \quad (8)$$

Note that

$$\sum_{i=\lfloor N/10 \rfloor}^N \frac{1}{i^2} = 9N^{-1} + \mathcal{O}(N^{-2}). \quad (9)$$

It remains to estimate

$$\sum_{i=0}^N \left(\sum_{j=0}^{n-1} ((i_0 10^{-j} - \lfloor i_0 10^{-j} \rfloor) - (i 10^{-j} - \lfloor i 10^{-j} \rfloor)) \right)^2,$$

which can be rewritten as

$$\begin{aligned} & \sum_{i=0}^N \left\{ \sum_{j=0}^{n-1} ((i_0 10^{-j} - \lfloor i_0 10^{-j} \rfloor) - (i 10^{-j} - \lfloor i 10^{-j} \rfloor)) \right\}^2 \\ & = (N+1) \left(\sum_{j=0}^{n-1} (i_0 10^{-j} - \lfloor i_0 10^{-j} \rfloor) \right)^2 + \sum_{i=0}^N \left(\sum_{j=0}^{n-1} (i 10^{-j} - \lfloor i 10^{-j} \rfloor) \right)^2 \\ & \quad - 2 \sum_{j=0}^{n-1} (i_0 10^{-j} - \lfloor i_0 10^{-j} \rfloor) \sum_{i=0}^N \sum_{j=0}^{n-1} (i 10^{-j} - \lfloor i 10^{-j} \rfloor). \end{aligned} \quad (10)$$

In the following, we consider each term on the right-hand side of (10) separately.

First we consider $\sum_{i=0}^N \sum_{j=0}^{n-1} (i 10^{-j} - \lfloor i 10^{-j} \rfloor)$, by switching the order of the summations. For every $i = 0, \dots, N$ and $j = 0, \dots, n-1$, there exist nonnegative integers k, l with $l \leq 10^j - 1$ such that $i = k 10^j + l$, and hence

$$i 10^{-j} - \lfloor i 10^{-j} \rfloor = l 10^{-j}.$$

Therefore,

$$\begin{aligned} & \{i : 0 \leq i \leq N\} \\ &= \{k10^j + l : 0 \leq k \leq \lfloor N10^{-j} \rfloor - 1, 0 \leq l \leq 10^j - 1\} \\ &\cup \{\lfloor N10^{-j} \rfloor 10^j + l : 0 \leq l \leq N - \lfloor N10^{-j} \rfloor 10^j\}. \end{aligned}$$

Let $N = \overline{a_{n-1} \cdots a_0} = \sum_{j=0}^{n-1} a_j 10^j$ with $0 \leq a_j \leq 9$ for all $0 \leq j \leq n-1$.

Notice that

$$\lfloor N10^{-j} \rfloor = N10^{-j} - \sum_{r=0}^{j-1} a_r 10^{r-j}, \quad \forall j = 1, \dots, n-1. \quad (11)$$

From the simple observations

$$\sum_{j=0}^{n-1} (1 - 10^{-j}) = n + \mathcal{O}(1)$$

and

$$\sum_{j=1}^{n-1} 10^{-j} \sum_{r=0}^{j-1} a_r 10^r = \mathcal{O}(n),$$

it is tedious but straightforward to deduce that

$$\begin{aligned} & \sum_{i=0}^N \sum_{j=0}^{n-1} (i10^{-j} - \lfloor i10^{-j} \rfloor) \\ &= \frac{1}{2} \left(n - \frac{10}{9} \right) N - \frac{1}{2} \sum_{j=1}^{n-1} \sum_{r=0}^{j-1} a_r 10^r + \frac{1}{2} \sum_{j=1}^{n-1} 10^{-j} \left(\sum_{r=0}^{j-1} a_r 10^r \right)^2 + \mathcal{O}(n). \end{aligned} \quad (12)$$

Next, we deal with $\sum_{i=0}^N \left(\sum_{j=0}^{n-1} (i10^{-j} - \lfloor i10^{-j} \rfloor) \right)^2$, which can be expanded as

$$\begin{aligned} & \sum_{i=0}^N \left(\sum_{j=0}^{n-1} (i10^{-j} - \lfloor i10^{-j} \rfloor) \right)^2 \\ &= 2 \sum_{i=0}^N \sum_{j=1}^{n-1} (i10^{-j} - \lfloor i10^{-j} \rfloor) \sum_{r=0}^{j-1} (i10^{-r} - \lfloor i10^{-r} \rfloor) + \sum_{i=0}^N \sum_{j=0}^{n-1} (i10^{-j} - \lfloor i10^{-j} \rfloor)^2. \end{aligned} \quad (13)$$

For every $1 \leq j \leq n-1$, let $K_j = N - \lfloor N10^{-j} \rfloor 10^j$ for notational convenience. Then similarly,

$$\begin{aligned} & \{i : 0 \leq i \leq N\} \\ &= \{k10^j + p10^r + l : 0 \leq k \leq \lfloor N10^{-j} \rfloor - 1, 0 \leq p \leq 10^{j-r} - 1, 0 \leq l \leq 10^r - 1\} \\ &\cup \{\lfloor N10^{-j} \rfloor 10^j + p10^r + l : 0 \leq p \leq \lfloor K_j 10^{-r} \rfloor - 1, 0 \leq l \leq 10^r - 1\} \\ &\cup \{\lfloor N10^{-j} \rfloor 10^j + \lfloor K_j 10^{-r} \rfloor 10^r + l : 0 \leq l \leq K_j - \lfloor K_j 10^{-r} \rfloor 10^r\}, \end{aligned}$$

from which it follows that

$$\begin{aligned} & \sum_{i=0}^N \sum_{j=1}^{n-1} (i10^{-j} - \lfloor i10^{-j} \rfloor) \sum_{r=0}^{j-1} (i10^{-r} - \lfloor i10^{-r} \rfloor) \\ &= \sum_{j=1}^{n-1} \sum_{r=0}^{j-1} \left(\sum_{k=0}^{\lfloor N10^{-j} \rfloor - 1} \sum_{p=0}^{10^{j-r}-1} \sum_{l=0}^{10^r-1} (p10^r + l) \cdot 10^{-j} l 10^{-r} \right. \\ &\quad + \sum_{p=0}^{\lfloor K_j 10^{-r} \rfloor - 1} \sum_{l=0}^{10^r-1} (p10^r + l) 10^{-j} l 10^{-r} \\ &\quad \left. + \sum_{l=0}^{K_j - \lfloor K_j 10^{-r} \rfloor 10^r} (\lfloor K_j 10^{-r} \rfloor 10^r + l) 10^{-j} l 10^{-r} \right). \end{aligned} \tag{14}$$

From (11) and (14), a lengthy but elementary calculation leads to

$$\begin{aligned} & \sum_{i=0}^N \sum_{j=0}^{n-1} (i10^{-j} - \lfloor i10^{-j} \rfloor) \sum_{r=0}^{j-1} (i10^{-r} - \lfloor i10^{-r} \rfloor) \\ &= \left(\frac{n^2}{8} - \frac{85n}{216} \right) N - \frac{1}{4} \sum_{j=1}^{n-1} j \sum_{l=0}^{j-1} a_l 10^l + \frac{1}{4} \sum_{j=1}^{n-1} j 10^{-j} \left(\sum_{l=0}^{j-1} a_l 10^l \right)^2 + \mathcal{O}(N). \end{aligned}$$

Analogously, one obtains also $\sum_{i=0}^N \sum_{j=0}^{n-1} (i10^{-j} - \lfloor i10^{-j} \rfloor)^2 = \frac{nN}{3} + \mathcal{O}(N)$. Note that (13) immediately leads to

$$\begin{aligned} & \sum_{i=0}^N \left(\sum_{j=0}^{n-1} (i10^{-j} - \lfloor i10^{-j} \rfloor) \right)^2 \\ &= \left(\frac{n^2}{4} - \frac{49}{108} n \right) N - \frac{1}{2} \sum_{j=0}^{n-1} j \sum_{l=0}^{j-1} a_l 10^l + \frac{1}{2} \sum_{j=0}^{n-1} j 10^{-j} \left(\sum_{l=0}^{j-1} a_l 10^l \right)^2 + \mathcal{O}(N). \end{aligned} \tag{15}$$

The rest of the proof consists of choosing an appropriate i_0 (or equivalently, $y = \langle \log_{10} i_0 \rangle$) to obtain a sufficiently precise bound for (10): Let

$$i_0 = 10^{n-1} - 10^{\lfloor n/2 \rfloor - 1} + 1 \quad \text{if} \quad 10^{n-1} \leq N \leq 10^n - 10^{\lfloor n/2 \rfloor},$$

and

$$i_0 = 10^n - 10^{\lfloor n/2 \rfloor} \quad \text{if} \quad 10^n - 10^{\lfloor n/2 \rfloor} < N \leq 10^n - 1.$$

Note that $\lfloor N/10 \rfloor + 1 \leq i_0 \leq N - 1$, and it is straightforward to verify that

$$\sum_{j=0}^{n-1} (i_0 10^{-j} - \lfloor i_0 10^{-j} \rfloor) = \frac{n}{2} + c + \mathcal{O}(N^{-1/2}) \quad \text{for some finite constant } c. \quad (16)$$

Combining (12), (15) and (16) yields

$$\begin{aligned} & \sum_{i=0}^N \left\{ \sum_{j=0}^{n-1} ((i 10^{-j} - \lfloor i 10^{-j} \rfloor) - (i_0 10^{-j} - \lfloor i_0 10^{-j} \rfloor)) \right\}^2 \\ &= \frac{11}{108} nN + \frac{1}{2} \sum_{j=0}^{n-1} (n-j) \left(\sum_{l=0}^{j-1} a_l 10^l \right) \left(1 - \sum_{l=0}^{j-1} a_l 10^{l-j} \right) + \mathcal{O}(N). \end{aligned}$$

Next, observe that

$$\frac{1}{2} \sum_{j=0}^{n-1} (n-j) \left(\sum_{l=0}^{j-1} a_l 10^l \right) \left(1 - \sum_{l=0}^{j-1} a_l 10^{l-j} \right) = \mathcal{O}(N),$$

which implies that

$$\begin{aligned} & \sum_{i=0}^N \left\{ \sum_{j=0}^{n-1} ((i 10^{-j} - \lfloor i 10^{-j} \rfloor) - (i_0 10^{-j} - \lfloor i_0 10^{-j} \rfloor)) \right\}^2 \\ &= \frac{11}{108} nN + \mathcal{O}(N), \end{aligned}$$

and hence

$$\begin{aligned} & \sum_{i=\lfloor N/10 \rfloor}^N \left\{ \sum_{j=0}^{n-1} ((i 10^{-j} - \lfloor i 10^{-j} \rfloor) - (i_0 10^{-j} - \lfloor i_0 10^{-j} \rfloor)) \right\}^2 \\ & \leq \frac{11 \log N}{108 \log 10} N + \mathcal{O}(N). \end{aligned} \quad (17)$$

Let $y = \langle \log_{10} i_0 \rangle$. Combining (7), (8), (9) and (17) yields

$$\begin{aligned} & \int_0^1 \left| (F_{\nu_N \circ \iota^{-1}}(x) - F_{\eta_N \circ \iota^{-1}}(x)) - (F_{\nu_N \circ \iota^{-1}}(y) - F_{\eta_N \circ \iota^{-1}}(y)) \right| dx \\ & \leq \frac{1}{6 \log 10} \sqrt{\frac{33}{\log 10}} \frac{\sqrt{\log N}}{N} + \mathcal{O}(N^{-1}); \end{aligned}$$

and hence with Proposition 2.1, it follows at long last that

$$\limsup_{N \rightarrow \infty} \frac{N}{\sqrt{\log N}} d_{\mathbb{T}}(\nu_N, \eta_N) \leq \frac{1}{6 \log 10} \sqrt{\frac{33}{\log 10}}.$$

□

Note that Theorem 3.1 describes the asymptotics of $(\nu_N(\log_b n))_{N \geq 1}$, in that it not only gives the rate of convergence, but also identifies the exponential distribution with specific rotation that (ν_N) asymptotically approaches.

REMARK 3.2. (i) It follows from a general result in [14] that

$$\limsup_{N \rightarrow \infty} \frac{N}{\log N} d_{\mathbb{T}}(\nu_N, E_{\log b} \circ R_{-\log_b N}^{-1}) < +\infty$$

for every $b \in \mathbb{N} \setminus \{1\}$. Obviously, this is weaker than (4).

(ii) From Zador's theorem on asymptotic quantization error in $\mathcal{P}(\mathbb{T})$ [4, Thm.1.4], it follows that

$$\liminf_{N \rightarrow \infty} N d_{\mathbb{T}}(\nu_N, E_{\log b} \circ R_{-\log_b N}^{-1}) > 0.$$

This shows that

$$\left(d_{\mathbb{T}}(\nu_N, E_{\log b} \circ R_{-\log_b N}^{-1}) \right)_{N \geq 1}$$

cannot decay faster than (N^{-1}) , and [1, Cor.3.8] suggests that it may be challenging to improve this lower bound.

(iii) Even if the inequality (8) is replaced by the following Hölder inequality,

$$\begin{aligned} \sum_{i=\lfloor N/10 \rfloor}^N \frac{1}{i} \left| \sum_{j=0}^{n-1} ((i_0 10^{-j} - \lfloor i_0 10^{-j} \rfloor) - (i 10^{-j} - \lfloor i 10^{-j} \rfloor)) \right| & \leq \left(\sum_{i=\lfloor N/10 \rfloor}^N \frac{1}{i^{4/3}} \right)^{3/4} \times \\ & \left(\sum_{i=\lfloor N/10 \rfloor}^N \left(\sum_{j=0}^{n-1} ((i_0 10^{-j} - \lfloor i_0 10^{-j} \rfloor) - (i 10^{-j} - \lfloor i 10^{-j} \rfloor)) \right)^4 \right)^{1/4}, \end{aligned}$$

the upper bound for the rate of convergence does not improve. Indeed, a tedious computation similar to the one in the proof of Theorem 3.1 yields

$$\limsup_{N \rightarrow \infty} \frac{N}{\sqrt{\log N}} d_{\mathbb{T}}(\nu_N, E_{\log b} \circ R_{-\log_b N}^{-1}) \leq c, \quad (18)$$

where the constant c may be smaller than $\frac{1}{6 \log 10} \sqrt{\frac{33}{\log 10}}$ but still is positive.

From this, one may optimistically conjecture that for all $b > 1$ (not necessarily integers), the sequence

$$\left(\frac{N}{\sqrt{\log N}} d_{\mathbb{T}}(\nu_N, E_{\log b} \circ R_{-\log_b N}^{-1}) \right)_{N \geq 2}$$

is bounded above and below by positive constants. Especially for non-integer b , this is speculation only, since many of the explicit calculations and estimates leading to (4) do not apply directly.

3.2. Sharp rates of convergence w.r.t. $d_{\mathbb{I}}$ and d_K

In this subsection, we complement the results of Subsection 3.1 by characterizing the sharp rate of convergence of $(\nu_N(\log_b n))_{N \geq 1}$ w.r.t. both $d_{\mathbb{I}}$ and d_K .

THEOREM 3.3. *Assume $b \in \mathbb{N} \setminus \{1\}$. Then*

$$\lim_{N \rightarrow \infty} \frac{N}{\log N} d_{\mathbb{I}}(\nu_N \circ \iota^{-1}, E_{\log b} \circ R_{-\log_b N}^{-1} \circ \iota^{-1}) = \frac{1}{2 \log b}.$$

Proof. Recall that $b = 10$. As in the proof of Theorem 3.1, by formula (2), it is easy to verify that for $10^{n-1} \leq N \leq 10^n - 1$,

$$\begin{aligned} & d_{\mathbb{I}}(\nu_N \circ \iota^{-1}, E_{\log b} \circ R_{-\log_b N}^{-1} \circ \iota^{-1}) \\ &= \frac{1}{N \log 10} \sum_{i=\lfloor N/10 \rfloor}^N \frac{1}{i} \sum_{j=0}^{n-1} (i10^{-j} - \lfloor i10^{-j} \rfloor) + \mathcal{O}(N^{-1}). \end{aligned} \quad (19)$$

Like the expression for

$$\sum_{i=0}^N \sum_{j=0}^{n-1} (i10^{-j} - \lfloor i10^{-j} \rfloor)$$

as in the proof of Theorem 3.3, it is readily checked that

$$\begin{aligned} \sum_{i=\lfloor N/10 \rfloor}^N \frac{1}{i} \sum_{j=0}^{n-1} (i10^{-j} - \lfloor i10^{-j} \rfloor) = \\ \sum_{j=0}^{n-1} \left(\sum_{l=\lfloor N/10 \rfloor - \lfloor \lfloor N/10 \rfloor 10^{-j} \rfloor 10^j}^{10^j-1} \frac{l10^{-j}}{\lfloor \lfloor N/10 \rfloor 10^{-j} \rfloor 10^j + l} + \right. \\ \left. \sum_{k=\lfloor \lfloor N/10 \rfloor 10^{-j} \rfloor + 1}^{\lfloor N10^{-j} \rfloor - 1} \sum_{l=0}^{10^j-1} \frac{l10^{-j}}{k10^j + l} + \sum_{l=0}^{N - \lfloor N10^{-j} \rfloor 10^j} \frac{l10^{-j}}{\lfloor N10^{-j} \rfloor 10^j + l} \right), \end{aligned}$$

which implies that

$$\begin{aligned} \sum_{i=\lfloor N/10 \rfloor}^N \frac{1}{i} \sum_{j=0}^{n-1} (i10^{-j} - \lfloor i10^{-j} \rfloor) \\ \geq \sum_{j=0}^{n-1} \left(\sum_{l=\lfloor N/10 \rfloor - \lfloor \lfloor N/10 \rfloor 10^{-j} \rfloor 10^j}^{10^j-1} \frac{l10^{-j}}{\lfloor \lfloor N/10 \rfloor 10^{-j} \rfloor 10^j + 10^j} \right. \\ \left. + \sum_{k=\lfloor \lfloor N/10 \rfloor 10^{-j} \rfloor + 1}^{\lfloor N10^{-j} \rfloor - 1} \sum_{l=0}^{10^j-1} \frac{l10^{-j}}{k10^j + 10^j} + \sum_{l=0}^{N - \lfloor N10^{-j} \rfloor 10^j} \frac{l10^{-j}}{N} \right) \\ = \frac{1}{2} \sum_{j=0}^{n-1} \left(\frac{1 - 10^{-j} + \lfloor N/10 \rfloor 10^{-j} - \lfloor \lfloor N/10 \rfloor 10^{-j} \rfloor}{\lfloor \lfloor N/10 \rfloor 10^{-j} \rfloor + 1} \right. \\ \times (1 - \lfloor N/10 \rfloor 10^{-j} + \lfloor \lfloor N/10 \rfloor 10^{-j} \rfloor) + \frac{N10^{-j} - \lfloor N10^{-j} \rfloor}{N} \\ \left. \times (N10^{-j} - \lfloor N10^{-j} \rfloor + 1) 10^j + (1 - 10^{-j}) \sum_{k=\lfloor \lfloor N/10 \rfloor 10^{-j} \rfloor + 1}^{\lfloor N10^{-j} \rfloor - 1} \frac{1}{k+1} \right). \end{aligned} \quad (20)$$

Note that

$$\begin{aligned} \frac{1}{2} \sum_{j=0}^{n-1} \left(\frac{1 - 10^{-j} + \lfloor N/10 \rfloor 10^{-j} - \lfloor \lfloor N/10 \rfloor 10^{-j} \rfloor}{\lfloor \lfloor N/10 \rfloor 10^{-j} \rfloor + 1} \right. \\ \times (1 - \lfloor N/10 \rfloor 10^{-j} + \lfloor \lfloor N/10 \rfloor 10^{-j} \rfloor) \\ \left. + \frac{(N10^{-j} - \lfloor N10^{-j} \rfloor)}{N} (N10^{-j} - \lfloor N10^{-j} \rfloor + 1) 10^j \right) = \mathcal{O}(1). \end{aligned}$$

Moreover, it is tedious but straightforward to confirm that

$$\frac{1}{2} \sum_{j=0}^{n-1} (1 - 10^{-j}) \sum_{k=\lfloor N/10 \rfloor 10^{-j} + 1}^{\lfloor N 10^{-j} \rfloor - 1} \frac{1}{k+1} = \frac{1}{2} \log N + \mathcal{O}(1),$$

with which (20) takes the form

$$\sum_{i=\lfloor N/10 \rfloor}^N \frac{1}{i} \sum_{j=0}^{n-1} (i 10^{-j} - \lfloor i 10^{-j} \rfloor) \geq \frac{1}{2} \log N + \mathcal{O}(1). \quad (21)$$

Analogously, one can also show that (21) holds with $\geq$ replaced by $\leq$, and hence

$$\sum_{i=\lfloor N/10 \rfloor}^N \frac{1}{i} \sum_{j=0}^{n-1} (i 10^{-j} - \lfloor i 10^{-j} \rfloor) = \frac{1}{2} \log N + \mathcal{O}(1).$$

The conclusion now follows from (19). $\square$

The following corollary is immediately obtained from Theorem 3.3, together with [9, Thm.5] and the fact that $d_{\mathbb{I}} \leq d_K$.

COROLLARY 3.4. *Assume $b \in \mathbb{N} \setminus \{1\}$. Then*

$$\begin{aligned} 0 &< \liminf_{N \rightarrow \infty} \frac{N}{\log N} d_K(\nu_N \circ \iota^{-1}, E_{\log b} \circ R_{-\log_b N}^{-1} \circ \iota^{-1}) \\ &\leq \limsup_{N \rightarrow \infty} \frac{N}{\log N} d_K(\nu_N \circ \iota^{-1}, E_{\log b} \circ R_{-\log_b N}^{-1} \circ \iota^{-1}) < +\infty. \end{aligned}$$

Comparing Theorems 3.1 and 3.3, as well as Corollary 3.4, notice how

$$\left(d_{\mathbb{I}}(\nu_N, E_{\log b} \circ R_{-\log_b N}^{-1}) \right)_{N \geq 1}$$

decays somewhat faster than both

$$\left(d_{\mathbb{I}}(\nu_N \circ \iota^{-1}, E_{\log b} \circ R_{-\log_b N}^{-1} \circ \iota^{-1}) \right)_{N \geq 1}$$

and

$$\left(d_K(\nu_N \circ \iota^{-1}, E_{\log b} \circ R_{-\log_b N}^{-1} \circ \iota^{-1}) \right)_{N \geq 1}.$$

Moreover, the ratio

$$\frac{d_{\mathbb{I}}(\nu_N \circ \iota^{-1}, E_{\log b} \circ R_{-\log_b N}^{-1} \circ \iota^{-1})}{d_K(\nu_N \circ \iota^{-1}, E_{\log b} \circ R_{-\log_b N}^{-1} \circ \iota^{-1})}$$

is bounded above and below by positive constants. This is remarkable since

$$\inf_{\mu \neq \nu, \mu, \nu \in \mathcal{P}(\mathbb{T})} \frac{d_{\mathbb{I}}(\mu \circ \iota^{-1}, \nu \circ \iota^{-1})}{d_K(\mu \circ \iota^{-1}, \nu \circ \iota^{-1})} = 0.$$

ACKNOWLEDGEMENT. Deepest thanks to his PhD thesis advisor Arno Berger for proposing this project, many helpful discussions, as well as the effort in improving the presentation of this manuscript as one chapter in the author's thesis. The author is also indebted to one anonymous referee for his/her serious work including pointing out a mistake in the proof of Theorem 3.1 as well as other valuable comments which also help improve the presentation of the current manuscript.

REFERENCES

- [1] CABRELLI, C. A.—MOLTER, U. M.: *The Kantorovich metric for probability measures on the circle*, J. Comput. Appl. Math. **57** (1995), no. 3, 345–361.
- [2] DUDLEY, R. M.: *Real Analysis and Probability*, 2nd ed., Cambridge Univ. Press, 2004.
- [3] GIULIANO ANTONINI, R.—STRAUCH, O.: *On weighed distribution functions of sequences*, Unif. Distrib. Theory **3** (2008), no. 1, 1–18.
- [4] IACOBELLI, M.: *Asymptotic quantization for probability measures on Riemannian manifolds*, ESAIM: Control Optim. Calc. Var. **22** (2016), no. 3, 770–785.
- [5] KEMPERMAN, J. H. B.: *Distributions modulo 1 of slowly changing sequences*, Nieuw Arch. Wisk. **21** (1973), no. 3, 138–163.
- [6] KUIPERS, L.—NIEDERREITER, H.: *Uniform Distribution of Sequences*, John Wiley & Sons, New York, 1974. Reprint edition: Dover Publications, Inc. Mineola, New York, 2006.
- [7] MARKLOF, J.—STRÖMBERGSSON, A.: *Gaps between logs*, Bull. London Math. Soc. **45** (2013), no. 6, 1267–1280.
- [8] OHKUBO, Y.: *On sequences involving primes*, Unif. Distrib. Theory **6** (2011), no. 2, 221–238.
- [9] OHKUBO, Y.—STRAUCH, O.: *Distribution of leading digits of numbers*, Unif. Distrib. Theory **11** (2016), no. 1, 23–45.
- [10] STRAUCH, O.—BLAŽEKOVÁ, O.: *Distribution of the sequence $p_n/n \bmod 1$* , Unif. Distrib. Theory, **1** (2006), no. 1, 45–63.
- [11] WINKLER, R.: *On the distribution behaviour of sequences*, Math. Nachr. **186** (1997), no. 1, 303–312.
- [12] WINTNER, A.: *On the cyclical distribution of the logarithms of the prime numbers*, Quart. J. Math. Oxford **6** (1935), no. 1, 65–66.
- [13] XU, C.: *On the rate of convergence for $(\log_b n)$* , preprint (2016), arXiv:1609.08207.
- [14] ———: *Rates of convergence for a class of slowly changing sequences* (in preparation).

CHUANG XU

Received October 2, 2017

Accepted September 12, 2018

Chuang Xu

Department of Mathematical Sciences

University of Copenhagen

Universitetsparken 5

DK-2100 Copenhagen

DENMARK

E-mail: chuang@math.ku.dk